



**U.S. GRAINS &
BIOPRODUCTS**
COUNCIL



**2025/2026
CORN EXPORT CARGO
QUALITY REPORT**



**U.S. GRAINS &
BIOPRODUCTS**
COUNCIL



Developing a report of this scope and breadth in a timely manner requires participation by several individuals and organizations. The U.S. Grains & BioProducts Council is grateful to Steve Hofing, Lee Singleton, Lisa Eckel, Dee Ann Hoffman and Erin Krannawitter of Centrec Consulting Group, LLC (Centrec) for their oversight and coordination in developing this report. A team of experts provided analysis and writing support. External team members include Drs. Lowell Hill, Marvin Paulsen and Tom Whitaker. In addition, the Council is indebted to the Illinois Crop Improvement Association's Identity Preserved Grain Laboratory (IPG Lab) and Champaign-Danville Grain Inspection (CDGI) for providing the corn quality testing services.

In particular, we acknowledge the irreplaceable services of the Federal Grain Inspection Service (FGIS) of the U.S. Department of Agriculture. FGIS provided samples from export cargoes. The FGIS Office of International Affairs coordinated the sampling process. FGIS field staff, the Washington State Department of Agriculture and FGIS-designated domestic official service providers collected and submitted the samples that constitute the foundation of this report. We are grateful for the time they devoted during their busy season.

As a USDA programs participant, the U.S. Grains & BioProducts Council is committed to complying with non-discrimination policies from federal, state and local civil rights laws and those of the USDA. Visit the USDA website page (<https://www.usda.gov/non-discrimination-statement>) for details.



1	Greetings from the Council	
2	Export Cargo Quality Highlights	
4	Introduction	
6	Quality Test Results	
	A. Grade Factors.....	6
	B. Chemical Composition.....	19
	C. Physical Factors	29
	D. Mycotoxins.....	48
59	U.S. Corn Export System	
	A. U.S. Corn Export Flow.....	60
	B. Impact of the Corn Marketing Channel on Quality.....	61
	C. U.S. Government Inspection and Grading.....	63
65	Survey & Statistical Analysis Methods	
	A. Overview.....	65
	B. Survey Design and Sampling	66
	C. Statistical Analysis	71
72	Testing Analysis Methods	
	A. Grade Factors.....	72
	B. Chemical Composition.....	73
	C. Physical Factors	74
	D. Mycotoxins.....	75
77	Historical Perspective	
85	U.S. Corn Supporting Information	
BC	USGBC Contact Information	



The U.S. Grains & BioProducts Council (USGBC) is pleased to present findings from its fifteenth annual corn quality survey in this *2025/2026 Corn Export Cargo Quality Report*.

The Council is committed to furthering global food security and mutual economic benefit through trade. In doing so, it offers this report to assist buyers in making well-informed decisions by providing reliable and timely information about the quality of U.S. corn destined for export.

The *Corn Export Cargo Quality Report* is the second of two reports released annually by the Council detailing the quality of the 2025 corn crop. The report is based on samples taken at the point of loading for international shipment early in the 2025/2026 marketing year. This report and its sister report, the *2025/2026 Corn Harvest Quality Report*, provide an early look at the grade factors established by the U.S. Department of Agriculture as well as chemical composition and other quality characteristics not reported elsewhere. This series of quality reports uses a consistent and transparent methodology to allow for insightful comparisons across time.

The Council is pleased to offer this report as a service to our valued trading partners. It serves as a means of fulfilling the Council's mission of developing markets, enabling trade and improving lives.



Sincerely,



Mark Wilson
Chairman, U.S. Grains & BioProducts Council
March 2026

The 2025 U.S. corn crop experienced generally favorable weather conditions throughout its development, which contributed to its projected record average yield and good grain quality. Reflecting this impact, the average aggregate quality of the corn samples tested for the *2025/2026 U.S. Grains & BioProducts Council Corn Export Cargo Quality Report (2025/2026 Export Cargo Report)* was better than or equal to U.S. No. 2 on all grade factors. Average broken corn and foreign material (BCFM) was the lowest observed in the fifteen years of the report, and total damage was lower than the 5YA¹. Additionally, 98.4% of samples tested below the U.S. Food and Drug Administration (FDA) action level for aflatoxin, and 99.4% of samples tested below the FDA's advisory level for deoxynivalenol (DON) or vomitoxin. The following are notable U.S. Aggregate quality attributes from the 2025/2026 export samples.

GRADE FACTORS

- Average **test weight** (57.8 pounds per bushel (lb/bu) or 74.4 kilograms per hectoliter (kg/hl)) was lower than 2024/2025 and the 5YA. However, all (100.0%) of the samples tested at or above the minimum for U.S. No. 2 grade indicating overall good quality.
- Average **BCFM** (2.4%) was lower than 2024/2025, the 5YA and the maximum limit for U.S. No. 2 grade. BCFM predictably increased from 0.3 to 2.4% as the crop moved from harvest through the marketing channel to export.
- Average **total damage** at export (1.6%) was lower than 2024/2025 and the 5YA. Most (93.0%) of the samples were at or below the limit for U.S. No. 1 grade, compared to 92.3% of the samples in 2024/2025.
- Average **heat damage** was 0.0%, the same as 2024/2025 and the 5YA, indicating good management of drying and storage of corn throughout the marketing channel.
- Using data obtained from the Federal Grain Inspection Service (FGIS) Export Grain Report, average **moisture** content in 2025/2026 (14.2%) was estimated to be lower than the 5YA (14.3%). Given incomplete data from the Southern Rail ECA, detailed moisture results are not presented in this *Export Cargo Report*. Further discussion of this protocol is included in the “Survey and Statistical Analysis Methods” section.

CHEMICAL COMPOSITION

- **Protein** concentration (8.5% dry basis) was similar to 2024/2025 but lower than the 5YA.
- **Starch** concentration (72.1% dry basis) was the same as 2024/2025 and higher than the 5YA.
- **Oil** concentration (3.8% dry basis) was lower than 2024/2025 and the 5YA.

¹The 5YA represents the simple average of the quality factors' average or standard deviation from the 2020/2021, 2021/2022, 2022/2023, 2023/2024 and 2024/2025 Export Cargo Reports.

PHYSICAL FACTORS

- Average **stress cracks** (15.0%) was higher than 2024/2025 and the 5YA. Of this year's export samples, 65.9% had less than 15% stress cracks compared to 70.8% of the samples in 2024/2025.
- Average **100-kernel weight** (36.83 grams) was similar to 2024/2025 and higher than the 5YA, indicating heavier kernels in 2025/2026 than the 5YA.
- Average **kernel volume** (0.29 cubic centimeters (cm³)) was the same as 2024/2025 and similar to the 5YA.
- Average **true density** (1.289 grams per cubic centimeter (g/cm³)) was lower than 2024/2025 but higher than the 5YA.
- Average percent of **whole kernels** (82.6%) was lower than 2024/2025 and the 5YA.
- Average **horneous (hard) endosperm** (84%) was lower than 2024/2025 but the same as the 5YA.


MYCOTOXINS

- All but three samples (98.4%) tested below the FDA action level of 20.0 parts per billion (ppb) for **afatoxin**. A total of 96.2% of the export samples had levels of aflatoxin below the FGIS "Lower Conformance Limit" of 5.0 ppb in 2025/2026, a similar proportion to 2024/2025 (96.1%) and higher than 2023/2024 (91.2%).
- All but one sample (99.4%) tested below the 5.0 parts per million (ppm) FDA advisory level for **DON**. Of the samples tested for DON in 2025/2026, 93.4% showed levels below 1.5 ppm, a lower proportion than 2024/2025 (96.7%) but the same as 2023/2024.
- Of the samples tested for **fumonisin** in 2025/2026, 96.7% were below the FDA's strictest guidance level for fumonisin of 5.0 ppm, a similar proportion to 2024/2025 (97.8%) and 2023/2024 (97.2%).
- This is the fifth year in which samples were tested for **ochratoxin A**, trichothecenes (**T-2**) and **zearalenone**. The "Quality Test Results" section contains sample results for these additional mycotoxins.

Corn quality information is important to foreign buyers as they decide on purchase contracts and processing needs for feed, food or industrial use. The *2025/2026 Export Cargo Report* provides accurate, unbiased information about the quality of U.S. yellow commodity corn as it is assembled for export early in the marketing year. This report provides test results for corn samples collected during the U.S. government-licensed sampling and inspection processes for U.S. corn waterborne and rail export shipments.

This *Export Cargo Report* is based on 430 yellow commodity corn samples collected from corn export shipments as they underwent the federal inspection and grading processes performed by FGIS or licensed inspectors at interior offices. The sample test results are reported at the U.S. aggregate level (U.S. Aggregate) and by export points associated with three general regions known as Export Catchment Areas (ECAs) shown on the map.

Export Catchment Areas



The grade factor results are also summarized by grade categories “U.S. No. 2” and “U.S. No. 3” to illustrate the practical quality differences between these two specifications. Results are displayed by the grade on the loading contract specified by the trading partners rather than their actual grade factor test results.

This report provides detailed information on each of the quality factors tested, including average, standard deviation and distribution, for the U.S. Aggregate and for each of the three ECAs. The “Quality Test Results” section summarizes the following quality factors:

- Grade Factors: test weight, BCFM, total damage and heat damage
- Chemical Composition: protein, starch and oil concentrations
- Physical Factors: stress cracks, 100-kernel weight, kernel volume, kernel true density, whole kernels and horneous (hard) endosperm
- Mycotoxins: aflatoxin, DON, fumonisin, ochratoxin A, T-2 and zearalenone

Details about the testing analysis methods used for this report are provided in the “Testing Analysis Methods” section.

For the *2025/2026 Export Cargo Report*, FGIS and interior offices collected samples from export shipments loaded from November 2025 through late-February 2026 to generate statistically valid results for the U.S. Aggregate and by ECA. The objective was to obtain enough samples to estimate quality factor averages of the corn exports with a relative margin of error (Relative ME) of not more than 10% for the U.S. Aggregate level.

Beginning with the *2022/2023 Export Cargo Report*, grade factor results from the Gulf and Pacific Northwest ECAs for the current year and all previous years were obtained from the Federal Grain Inspection Service's Export Grain Report. This dataset provides shipment-level grade factor and moisture data, the contract grade specified by the trading partners, the metric tons shipped and the inspection date. These data were filtered by date to correspond with the timeframe from which the physical samples were collected using the dataset's Thursday cutoff date. Complete grade factor data are available from nearly all vessels loaded from the Gulf and Pacific Northwest ECAs, which provide the best possible representation of the average grade factors loaded from these ECAs since all vessels loaded with corn are sampled.¹ While the dataset contains shipment information from the Southern Rail ECA, this report continues to rely on the grade factor results from the physical samples collected since grade factor data are often incomplete from rail shipments in the FGIS dataset.² Further discussion of this protocol and other statistical sampling and analysis methods are presented in the "Survey and Statistical Analysis Methods" section. Annual inspection data are available to the public at <https://fgisonline.ams.usda.gov/ExportGrainReport/default.aspx>.

This *2025/2026 Export Cargo Report* is the fifteenth in a series of annual surveys of the quality of U.S. corn exports early in the marketing year. In addition to the Council reporting the quality of corn exports early in the current marketing year, the cumulative *Export Cargo Report* surveys are providing increased value to stakeholders. The fifteen years of data enable export buyers and other stakeholders to make year-to-year comparisons and assess patterns in corn quality based on growing, drying, handling, storage and transport conditions.

The *Export Cargo Report* does not predict the actual quality of any cargo or lot of corn after loading or at destination. All participants in the value chain need to understand their own contract needs and obligations. In addition to grade, many of the quality attributes can be specified in the buyer-seller contract. Many factors, including weather, genetics, commingling and grain drying and handling, affect quality changes in complex ways. Sample test results can vary significantly depending on the origination of the corn, the ways in which a corn lot was loaded onto a conveyance and the method of sampling used. A review of how corn quality evolves from the field to the ocean vessel or railcar is provided in the "U.S. Corn Export System" section.

The companion report, the *U.S. Grains & BioProducts Council 2025/2026 Corn Harvest Quality Report*, was released in November 2025 and reported on the quality of the corn as it entered the U.S. marketing system. The *2025/2026 Harvest Report* and the *2025/2026 Export Cargo Report* should be studied together so that changes in corn quality occurring between harvest and export can be understood. The "Historical Perspective" section illustrates these changes by displaying the results from this report with all previous *Harvest Reports* and *Export Cargo Reports*.

¹While this protocol results in a sampling population containing nearly all of the corn exported through the Gulf and Pacific Northwest ECAs, there are a few exclusions discussed in the "Survey and Statistical Analysis Methods" section.

²While often incomplete, grade factor and moisture data from rail shipments to Mexico were collected from the FGIS dataset to permit the estimation of U.S. Aggregate average moisture content. Detailed moisture results are not presented in this *Export Cargo Report* due to incomplete data from the Southern Rail ECA. However, a note is included on page two in the "Highlights" section indicating the U.S. Aggregate average moisture content derived from the FGIS data.

A. GRADE FACTORS

USDA FGIS has established numerical grades, definitions and standards for the measurement of many quality attributes. The attributes that determine the numerical grades for corn are test weight, BCFM, total damage and heat damage. A table with the numerical requirements for these attributes is in the “U.S. Corn Supporting Information” section of this report and on the following page.

SUMMARY: GRADE FACTORS

- Average U.S. Aggregate test weight (57.8 lb/bu or 74.4 kg/hl) was lower than 2024/2025 (58.3 lb/bu), 2023/2024 (58.1 lb/bu) and the 5YA (58.0 lb/bu) but higher than the 10YA (57.6 lb/bu). It was also well above the minimum for U.S. No. 1 grade (56.0 lb/bu).
- The average test weight at export has been consistently lower than that at harvest.
- The average test weight for the Pacific Northwest ECA has been consistently lower than that of the other ECAs over the last five years, the 5YA and the 10YA.
- Average U.S. Aggregate BCFM (2.4%) was lower than 2024/2025 and 2023/2024 (both 2.5%), the 5YA (2.6%) and the 10YA (2.8%). It was also below the U.S. No. 2 grade limit (3.0%).
- A total of 89.9% of the export samples was at or below the maximum BCFM allowed for U.S. No. 2 grade (3.0%) and 99.8% were at or below the BCFM limit for U.S. No. 3 grade (4.0%).
- Average BCFM in the Southern Rail ECA (1.1%) was lower than the Gulf (2.6%) and Pacific Northwest ECAs (3.1%). Average BCFM for the Southern Rail ECA has also been lowest among the ECAs for the last five years, the 5YA and the 10YA.

- Average U.S. Aggregate total damage (1.6%) was lower than 2024/2025 (1.8%), 2023/2024 (1.7%), the 5YA (2.0%) and the 10YA (2.2%). It was also well below the limit for U.S. No. 1 grade (3.0%).
- Of the export samples, 93.0% had 3.0% or less damaged kernels, meeting the U.S. No. 1 grade. In addition, 97.6% were at or below the limit for U.S. No. 2 grade (5.0%). Average total damage for the Pacific Northwest ECA has been the lowest among the ECAs for the last five years, the 5YA and the 10YA.
- Average U.S. Aggregate heat damage was 0.0% for 2025/2026, the same as the previous five years, the 5YA and 10YA.

U.S. Corn Grades and Grade Requirements				
Grade	Minimum Test Weight per Bushel (Pounds)	Maximum Limits of		
		Damaged Kernels		Broken Corn and Foreign Material (Percent)
		Heat Damaged (Percent)	Total (Percent)	
U.S. No. 1	56.0	0.1	3.0	2.0
U.S. No. 2	54.0	0.2	5.0	3.0
U.S. No. 3	52.0	0.5	7.0	4.0
U.S. No. 4	49.0	1.0	10.0	5.0
U.S. No. 5	46.0	3.0	15.0	7.0



TEST WEIGHT


Test weight (weight per volume) is a measure of bulk density. It is often used as a general indicator of overall quality and as a gauge of endosperm hardness for alkaline cookers and dry millers. High test weight corn takes up less storage space than the same weight of corn with lower test weight. Test weight is initially impacted by genetic differences in the structure of the kernel. However, it is also affected by moisture content, drying speed, physical damage to the kernel (broken or scuffed kernel surfaces), foreign material in the sample, kernel size and hardness, kernel maturity and microbiological damage. When sampled and measured at the point of delivery from the farm at a given moisture content, high test weight generally indicates high quality, a high percent of horneous (or hard) endosperm and sound, clean corn. Test weight has a positive correlation with true density and reflects kernel hardness and good maturation conditions.

RESULTS


- Average U.S. Aggregate test weight (57.8 lb/bu or 74.4 kg/hl) was well above the limit for U.S. No. 1 grade (56.0 lb/bu). It was also lower than 2024/2025 (58.3 lb/bu), 2023/2024 (58.1 lb/bu) and the 5YA (58.0 lb/bu) but higher than the 10YA (57.6 lb/bu).
- The 2025/2026 export samples had a standard deviation (0.67 lb/bu) similar to 2024/2025 (0.62 lb/bu), 2023/2024 (0.71 lb/bu), the 5YA (0.63 lb/bu) and the 10YA (0.68 lb/bu).

TEST WEIGHT (lb/bu & kg/hl)

U.S. Aggregate Results Summary




- Average U.S. Aggregate test weight at export (57.8 lb/bu) was lower than the 2025 harvest (58.6 lb/bu or 75.4 kg/hl). The average test weight at export has been lower than at harvest in each of the past fifteen years, with an average decrease from harvest to export of 0.6 lb/bu or 0.8 kg/hl.
- The variability of the test weight in the 2025/2026 export samples as measured by the standard deviation (0.67 lb/bu) was less than that of the 2025 harvest samples (1.37 lb/bu). As corn is commingled moving through the marketing channel, test weight becomes more uniform, with a lower standard deviation and a range between maximum and minimum values that is less than at harvest.
- The average test weight for the Pacific Northwest ECA (56.0 lb/bu) was lower than for the Southern Rail (57.4 lb/bu) and the Gulf ECAs (58.7 lb/bu). The average test weight for the Pacific Northwest ECA has consistently been lower than the other two ECAs over the last five years, the 5YA and the 10YA.
- U.S. Aggregate test weight in samples loaded as Grade No. 2 or No. 2 or better averaged 57.8 lb/bu while those loaded as Grade No. 3 or No. 3 or better averaged 57.7 lb/bu.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26




BROKEN CORN AND FOREIGN MATERIAL

BCFM is an indicator of the amount of clean, sound corn available for feeding and processing. The lower the percentage of BCFM, the less foreign material or fewer broken kernels are in the sample. As corn moves from farm deliveries through the marketing channel, each impact on the grain during handling and transporting increases the amount of broken corn. As a result, the average BCFM in most shipments of corn will be higher at the export point than in deliveries from the farm to the local elevator.



Broken corn (BC) is defined as corn and any other material (such as weed seeds) small enough to pass through a 12/64th-inch round-hole sieve but too large to pass through a 6/64th-inch round-hole sieve. Foreign material (FM) is defined as any non-corn material too large to pass through a 12/64th-inch round-hole sieve, as well as all fine material small enough to pass through a 6/64th-inch round-hole sieve.

RESULTS

- Average U.S. Aggregate BCFM in export samples (2.4%) was lower than 2024/2025 and 2023/2024 (both 2.5%), the 5YA (2.6%) and the 10YA (2.8%). It was also below the U.S. No. 2 grade limit (3.0%).
- The variability of the 2025/2026 export samples (with a standard deviation of 0.56%) was similar to 2024/2025 (0.60%), 2023/2024 (0.61%) and the 5YA (0.54%) and the same as the 10YA. The range in values (4.4%) was intermediate to 2024/2025 (4.9%) and 2023/2024 (3.8%).
- BCFM in the 2025/2026 export samples was distributed with 89.9% of the samples at or below the limit for U.S. No. 2 grade (3.0%), and 99.8% at or below the limit for U.S. No. 3 grade (4.0%).




- Average U.S. Aggregate BCFM at export (2.4%) was 2.1 percentage points higher than at harvest (0.3%). Both the 5YA and 10YA had a similar 1.9 to 2.0 percentage point increase between harvest and export. This increase in BCFM at export is likely due to higher broken corn as a result of artificial drying and additional impacts caused by conveying and handling equipment as the corn moves through the market channel.
- Average BCFM in the Southern Rail ECA (1.1%) was lower than the Gulf (2.6%) and Pacific Northwest ECAs (3.1%). Average BCFM for the Southern Rail has also been lowest among the ECAs for the previous five years, the 5YA and the 10YA.
- U.S. Aggregate BCFM in samples loaded as Grade No. 2 or No. 2 or better averaged 2.4% while those loaded as Grade No. 3 or No. 3 or better averaged 2.6%.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26



TOTAL DAMAGE

Total damage is the percent of kernels and pieces of kernels that are visually damaged in some way, including damage from mold, frost, insects, sprouting, disease, weather, ground, germ and heat. Most of these types of damage result in some discoloration or change in kernel texture. However, damage does not include broken pieces of grain that are otherwise normal in appearance.


Mold damage is usually associated with higher moisture content and warm temperatures during the growing season or during storage. Several field molds, such as *Diplodia*, *Aspergillus*, *Fusarium* and *Gibberella*, can lead to mold-damaged kernels during the growing season if the weather conditions are conducive to their development. While some fungi that produce mold damage can also produce mycotoxins, not all fungi produce mycotoxins. The chance of mold decreases as corn is dried and cooled to lower temperatures.

RESULTS


- Average U.S. Aggregate total damage (1.6%) was lower than 2024/2025 (1.8%), 2023/2024 (1.7%), the 5YA (2.0%) and the 10YA (2.2%). It was also well below the limit for U.S. No. 1 grade corn (3.0%).
- Variability in the 2025/2026 samples, as indicated by the standard deviation (0.83%), was higher than 2024/2025 (0.68%), similar to 2023/2024 (0.86%) and the 10YA (0.88%) but lower than the 5YA (1.02%). The 2025/2026 sample range (0.1 to 11.3%) was higher than 2024/2025 (0.2 to 7.9%) but similar to 2023/2024 (0.1 to 11.1%).

U.S. Aggregate Results Summary

TOTAL DAMAGE (%)




- Of the export samples, 93.0% had 3.0% or less damaged kernels, meeting the U.S. No. 1 grade. In addition, 97.6% were at or below the limit for U.S. No. 2 grade (5.0%).
- The average level of total damage in the marketing channel at export (1.6%) was higher than at harvest (1.1%). Total damage typically increases between harvest and export. The export 5YA (2.0%) was 1.0 percentage point higher than the harvest 5YA (1.0%); and the export 10YA (2.2%) was 0.7 percentage points higher than the harvest 10YA (1.5%). Total damage can increase during storage, especially if there are spout-lines and pockets of high moisture corn in storage bins or transport containers.
- Average total damage in the Pacific Northwest ECA (0.7%) was lower than the Gulf (1.7%) and Southern Rail ECAs (2.6%). Average total damage for the Pacific Northwest has been lowest among the ECAs for the last five years, the 5YA and the 10YA.
- U.S. Aggregate total damage in samples loaded as Grade No. 2 or No. 2 or better averaged 1.6% while those loaded as Grade No. 3 or No. 3 or better averaged 2.3%.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26





HEAT DAMAGE

Heat damage is a subset of total damage in corn grades and has separate allowances in the U.S. grade standards. Heat damage can be caused by microbiological activity in warm, moist grain or by high heat applied during drying. Low levels of heat damage may indicate the corn has been dried and stored at moisture contents and temperatures that prevent damage in the marketing channel.

RESULTS

- Average U.S. Aggregate heat damage for 2025/2026 was 0.0%, the same as 2024/2025, 2023/2024, the 5YA and 10YA (all 0.0%).
- These averages have been below the limit for U.S. No. 1 grade (0.1%), indicating good management of drying and storage of the corn throughout the marketing channel.

SUMMARY: GRADE FACTORS

2025/2026 Export Cargo						2024/2025 Export Cargo					2023/2024 Export Cargo				
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.
U.S. Aggregate						U.S. Aggregate					U.S. Aggregate				
Test Weight (lb/bu)	459	57.8	0.67	54.5	60.9	393	58.3*	0.62	56.1	60.3	381	58.1*	0.71	54.8	60.8
Test Weight (kg/hl)	459	74.4	0.86	70.2	78.3	393	75.1*	0.80	72.2	77.6	381	74.8*	0.91	70.5	78.2
BCFM (%)	459	2.4	0.56	0.3	4.7	430	2.5*	0.60	0.6	5.5	389	2.5*	0.61	0.3	4.1
Total Damage (%)	459	1.6	0.83	0.1	11.3	430	1.8*	0.68	0.2	7.9	389	1.7	0.86	0.1	11.1
Heat Damage (%)	459	0.0	0.00	0.0	0.1	430	0.0*	0.01	0.0	0.1	389	0.0	0.01	0.0	0.2
Gulf						Gulf					Gulf				
Test Weight (lb/bu)	282	58.7	0.58	57.2	60.9	288	58.7	0.51	57.0	60.3	239	58.9*	0.58	56.2	60.8
Test Weight (kg/hl)	282	75.6	0.75	73.6	78.3	288	75.6	0.66	73.4	77.6	239	75.8*	0.75	72.4	78.2
BCFM (%)	282	2.6	0.55	0.7	4.7	288	2.7*	0.50	0.7	4.7	239	2.8*	0.62	0.7	4.1
Total Damage (%)	282	1.7	0.56	0.7	4.7	288	2.1*	0.62	0.8	4.0	239	2.1*	0.75	0.4	4.6
Heat Damage (%)	282	0.0	0.01	0.0	0.1	288	0.0*	0.01	0.0	0.1	239	0.0	0.01	0.0	0.1
Pacific Northwest						Pacific Northwest					Pacific Northwest				
Test Weight (lb/bu)	120	56.0	0.73	54.5	57.7	91	57.1*	0.54	56.4	58.6	65	56.2*	0.76	54.8	58.7
Test Weight (kg/hl)	120	72.0	0.94	70.2	74.3	91	73.6*	0.70	72.6	75.5	65	72.4*	0.98	70.5	75.6
BCFM (%)	120	3.1	0.55	2.0	4.0	91	2.8*	0.68	0.9	4.0	65	3.2	0.56	2.1	4.0
Total Damage (%) ¹	120	0.7	0.31	0.1	1.6	91	0.7	0.29	0.2	1.6	65	0.6	0.41	0.1	2.2
Heat Damage (%)	120	0.0	0.00	0.0	0.0	91	0.0	0.01	0.0	0.1	65	0.0	0.01	0.0	0.1
Southern Rail						Southern Rail					Southern Rail				
Test Weight (lb/bu)	57	57.4	0.87	55.6	59.2	48	58.2*	0.99	56.1	59.7	77	57.7*	1.04	55.4	60.1
Test Weight (kg/hl)	57	73.9	1.12	71.6	76.2	48	74.9*	1.28	72.2	76.8	77	74.3*	1.34	71.3	77.4
BCFM (%) ¹	57	1.1	0.58	0.3	2.7	85	1.7*	0.83	0.6	5.5	85	0.9	0.63	0.3	3.9
Total Damage (%) ¹	57	2.6	2.19	0.2	11.3	85	2.1	1.27	0.4	7.9	85	1.6*	1.67	0.1	11.1
Heat Damage (%)	57	0.0	0.00	0.0	0.0	85	0.0	0.00	0.0	0.0	85	0.0	0.03	0.0	0.2

*Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

¹The relative margin of error for predicting the population average exceeded 10.0%.



SUMMARY: GRADE FACTORS

	Five-Year Average (MY20/21-MY24/25)			Ten-Year Average (MY15/16-MY24/25)		
	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.
U.S. Aggregate						
Test Weight (lb/bu)	1,953	58.0*	0.63	3,881	57.6*	0.68
Test Weight (kg/hl)	1,953	74.6*	0.81	3,881	74.2*	0.87
BCFM (%)	2,001	2.6*	0.54	3,929	2.8*	0.56
Total Damage (%)	2,001	2.0*	1.02	3,929	2.2*	0.88
Heat Damage (%)	2,001	0.0*	0.01	3,929	0.0*	0.01
Gulf						
Test Weight (lb/bu)	1,327	58.6*	0.57	2,732	58.2*	0.62
Test Weight (kg/hl)	1,327	75.4*	0.73	2,732	74.9*	0.80
BCFM (%)	1,327	2.8*	0.53	2,732	2.8*	0.53
Total Damage (%)	1,327	2.4*	0.87	2,732	2.6*	0.84
Heat Damage (%)	1,327	0.0	0.02	2,732	0.0*	0.02
Pacific Northwest						
Test Weight (lb/bu)	299	56.6*	0.57	513	56.2*	0.68
Test Weight (kg/hl)	299	72.9*	0.74	513	72.4*	0.87
BCFM (%)	299	3.0	0.51	513	3.3*	0.57
Total Damage (%)	299	0.8*	0.39	513	0.8*	0.41
Heat Damage (%)	299	0.0*	0.01	513	0.0*	0.01
Southern Rail						
Test Weight (lb/bu)	361	57.8*	0.90	670	57.7*	0.87
Test Weight (kg/hl)	361	74.4*	1.16	670	74.2*	1.12
BCFM (%)	409	1.7*	0.65	718	1.9*	0.62
Total Damage (%)	409	2.5	2.34	718	2.5	1.62
Heat Damage (%)	409	0.0*	0.01	718	0.0*	0.00

*Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

SUMMARY: GRADE FACTORS

2025/2026 Export Cargo Samples U.S. No. 2						2025/2026 Export Cargo Samples U.S. No. 3						2025 Harvest					
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples ¹	Avg.	Std. Dev.	Min.	Max.		
U.S. Aggregate						U.S. Aggregate						U.S. Aggregate					
Test Weight (lb/bu)	318	57.8	0.64	54.5	60.1	89	57.7	0.67	54.6	60.0	599	58.6**	1.37	52.6	63.6		
Test Weight (kg/hl)	318	74.3	0.83	70.2	77.3	89	74.3	0.87	70.3	77.2	599	75.4**	1.77	67.7	81.9		
BCFM (%)	318	2.4	0.43	0.8	3.0	89	2.6	0.85	0.8	4.7	599	0.3**	0.21	0.0	2.6		
Total Damage (%)	318	1.6	0.65	0.2	4.7	89	2.3*	0.63	0.1	6.5	599	1.1**	0.89	0.0	11.4		
Heat Damage (%)	318	0.0	0.00	0.0	0.1	89	0.0	0.00	0.0	0.0	599	0.0	0.00	0.0	0.0		
Gulf						Gulf						Gulf					
Test Weight (lb/bu)	239	58.7	0.54	57.2	60.1	29	58.5	0.73	57.2	60.0	572	58.7	1.50	52.6	63.6		
Test Weight (kg/hl)	239	75.6	0.70	73.6	77.3	29	75.3	0.94	73.7	77.2	572	75.6	1.93	67.7	81.9		
BCFM (%)	239	2.6	0.42	0.8	3.0	29	2.7	1.01	0.8	4.7	572	0.3**	0.21	0.0	2.6		
Total Damage (%)	239	1.7	0.53	0.7	4.7	29	1.7	0.71	0.8	4.0	572	1.1**	0.95	0.0	11.4		
Heat Damage (%)	239	0.0	0.01	0.0	0.1	29	0.0	0.00	0.0	0.0	572	0.0	0.00	0.0	0.0		
Pacific Northwest						Pacific Northwest						Pacific Northwest					
Test Weight (lb/bu)	64	56.1	0.69	54.5	57.6	56	55.8*	0.73	54.6	57.7	262	57.7**	1.13	52.9	62.3		
Test Weight (kg/hl)	64	72.2	0.89	70.2	74.1	56	71.8*	0.94	70.3	74.3	262	74.3**	1.46	68.1	80.2		
BCFM (%)	64	2.8	0.28	2.0	3.0	56	3.5*	0.54	2.1	4.0	262	0.4**	0.22	0.0	1.4		
Total Damage (%)	64	0.7	0.30	0.2	1.6	56	0.6	0.33	0.1	1.6	262	0.8**	0.66	0.0	7.4		
Heat Damage (%)	64	0.0	0.00	0.0	0.0	56	0.0	0.00	0.0	0.0	262	0.0	0.00	0.0	0.0		
Southern Rail						Southern Rail						Southern Rail					
Test Weight (lb/bu)	15	56.8	0.88	56.4	59.9	4	57.6*	0.44	56.9	58.4	342	59.3**	1.29	52.6	63.6		
Test Weight (kg/hl)	15	73.1	1.13	72.6	77.1	4	74.1*	0.57	73.2	75.2	342	76.3**	1.66	67.7	81.9		
BCFM (%)	15	1.6	0.62	1.6	3.0	4	1.2	0.70	2.2	3.1	342	0.3**	0.20	0.0	2.1		
Total Damage (%)	15	2.6	1.41	0.3	4.6	4	5.9*	0.73	0.3	6.5	342	1.4**	1.01	0.0	7.0		
Heat Damage (%)	15	0.0	0.00	0.0	0.0	4	0.0	0.00	0.0	0.0	342	0.0	0.00	0.0	0.0		






*Indicates average was significantly different from the average of samples loaded as U.S. No. 2 or better from this year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

**Indicates current year's Export Cargo average was significantly different from this year's Harvest, based on a 2-tailed t-test at the 95% level of confidence.

¹Due to the ECA results being composite statistics, the sum of the sample numbers from the three ECAs is greater than the U.S. Aggregate.



**GRADE FACTORS
AGGREGATE SIX-YEAR COMPARISON**



B. CHEMICAL COMPOSITION

The chemical composition of corn consists primarily of protein, starch and oil. While these attributes are not grade factors, they are of significant interest to end-users. Chemical composition values provide additional information related to nutritional value for livestock and poultry feeding, for wet milling uses and other processing uses of corn. Unlike many physical attributes, chemical composition values are not expected to change significantly during storage or transit.

SUMMARY: CHEMICAL COMPOSITION

- Average U.S. Aggregate protein concentration at export (8.5%) was similar to 2024/2025 (8.6%), lower than 2023/2024 (8.9%) and the 5YA (8.6%) but statistically different (higher) from the 10YA (8.5%).
- Average U.S. Aggregate starch concentration (72.1%) was the same as 2024/2025, higher than 2023/2024 (71.8%) and the 5YA (72.0%) but lower than the 10YA (72.3%). This year's average was also lower than the average U.S. Aggregate concentration from the 2025 harvest samples (72.3%).
- Average U.S. Aggregate oil concentration was 3.8% in 2025/2026. This value was lower than 2024/2025, 2023/2024, the 5YA and the 10YA (all 3.9%).
- The standard deviations for protein (0.34%), starch (0.38%) and oil (0.13%) in the export samples were lower than the standard deviations for protein (0.53%), starch (0.55%) and oil (0.23%) in the harvest samples.

PROTEIN


Protein is very important for poultry and livestock feeding because it supplies essential sulfur-containing amino acids and helps to improve feed conversion efficiency. Protein concentration tends to decrease with decreased available soil nitrogen and in years with high yields. On a single sample basis, protein is usually inversely related to starch concentration. Results are reported on a dry basis.

RESULTS


- Average U.S. Aggregate protein concentration in 2025/2026 (8.5%) was similar to 2024/2025 (8.6%), lower than 2023/2024 (8.9%) and the 5YA (8.6%) but statistically different (higher) from the 10YA (8.5%).
- Variability in the 2025/2026 samples, as indicated by the standard deviation (0.34%), was similar to 2024/2025 (0.35%), 2023/2024 (0.33%) and the 10YA (0.32%) and the same as the 5YA.
- The average protein concentration of the 2025/2026 export samples (8.5%) was higher than the average of the 2025 harvest samples (8.4%).

U.S. Aggregate Results Summary

PROTEIN (DRY BASIS %)




- The 2025/2026 export samples (standard deviation of 0.34%) was more uniform than the 2025 harvest samples (standard deviation of 0.53%). In addition, the range of protein concentrations at export (7.4 to 10.0%) was less than at harvest (5.4 to 11.1%). The uniformity is due, in part, to grains becoming more homogenous as they are aggregated from numerous harvest-level sources.
- The 2025/2026 export samples were distributed with 11.1% at or above 9.0%, compared with 14.4% of the 2024/2025 samples and 37.8% of the 2023/2024 samples.
- The Southern Rail and Gulf ECAs had the highest average protein concentration (8.6%), followed by the Pacific Northwest ECA (8.4%).



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26





STARCH


Starch is an important factor for corn used by wet millers and dry-grind ethanol manufacturers. High starch concentration is often indicative of good kernel growing/filling conditions and reasonably moderate kernel densities. Starch is usually inversely related to protein concentration on a single sample basis. Results are reported on a dry basis.

RESULTS


- Average U.S. Aggregate starch concentration (72.1%) was the same as 2024/2025, higher than 2023/2024 (71.8%) and the 5YA (72.0%) but lower than the 10YA (72.3%). This year's average was also lower than the average U.S. Aggregate concentration from the 2025 harvest samples (72.3%).
- Variability in the 2025/2026 samples, as indicated by the standard deviation (0.38%), was the same as 2024/2025 and 2023/2024 but similar to the 5YA (0.40%) and the 10YA (0.41%). The 2025/2026 sample range (70.8 to 73.4%) was similar to the range for 2024/2025 (70.8 to 73.3%) and the range for 2023/2024 (70.1 to 72.8%).

STARCH (DRY BASIS %)

U.S. Aggregate Results Summary




- The standard deviation for starch concentration of the 2025/2026 export samples (0.38%) was lower than the standard deviation of the 2025 harvest samples (0.55%). Average variability in export samples, as indicated by the standard deviation, is typically less at export than at harvest.
- Starch concentrations were distributed with 71.0% at or above 72.0%, compared to 68.0% in 2024/2025 and 32.1% in 2023/2024.
- Average starch concentration for the Pacific Northwest ECA (72.1%) was the same as the Southern Rail ECA and similar to that for the Gulf (72.2%).



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26





OIL


Oil is an essential component of poultry and livestock rations. It serves as an energy source, enables fat-soluble vitamins to be utilized and provides certain essential fatty acids. Oil is also an important co-product of corn wet and dry milling. Results are reported on a dry basis.

RESULTS


- Average U.S. Aggregate oil concentration was 3.8% in 2025/2026. This value was lower than 2024/2025, 2023/2024, the 5YA and the 10YA (all 3.9%).
- The 2025/2026 export samples had a standard deviation (0.13%) similar to 2024/2025 (0.12%) and 2023/2024 (0.14%).
- Average oil concentration for the 2025/2026 export samples (3.8%) was the same as the 2025 harvest samples (3.8%). The standard deviation at export (0.13%) was lower than at harvest (0.23%). Average oil concentration generally changes little from harvest to export, while variability in export samples is typically less at export than at harvest, as indicated by the lower standard deviation.

OIL (DRY BASIS %)

U.S. Aggregate Results Summary




- The 2025/2026 samples distribution showed 20.2% of samples at or above 4.0% oil as compared to 2024/2025 (25.4%) and 2023/2024 (27.2%).
- Average oil concentration for the Pacific Northwest ECA (3.9%) was higher than for the Gulf and Southern Rail ECAs (both 3.8%).
- There has been a 0.0 to 0.2% percentage point difference in average oil concentrations among the ECAs for each of the last five years, the 5YA and the 10YA.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26





SUMMARY: CHEMICAL COMPOSITION

2025/2026 Export Cargo						2024/2025 Export Cargo					2023/2024 Export Cargo				
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.
U.S. Aggregate						U.S. Aggregate					U.S. Aggregate				
Protein (Dry Basis %)	430	8.5	0.34	7.4	10.0	425	8.6	0.35	7.3	9.9	433	8.9*	0.33	8.0	10.0
Starch (Dry Basis %)	430	72.1	0.38	70.8	73.4	425	72.1	0.38	70.8	73.3	433	71.8*	0.38	70.1	72.8
Oil (Dry Basis %)	430	3.8	0.13	3.5	4.3	425	3.9*	0.12	3.5	4.2	433	3.9*	0.14	3.5	4.4
Gulf						Gulf					Gulf				
Protein (Dry Basis %)	248	8.6	0.27	7.7	9.5	247	8.6	0.27	7.9	9.5	246	8.9*	0.30	8.0	9.7
Starch (Dry Basis %)	248	72.2	0.33	71.0	73.1	247	72.1*	0.33	71.0	73.0	246	71.8*	0.35	70.1	72.6
Oil (Dry Basis %)	248	3.8	0.13	3.6	4.3	247	3.9	0.12	3.5	4.2	246	3.9*	0.13	3.6	4.4
Pacific Northwest						Pacific Northwest					Pacific Northwest				
Protein (Dry Basis %)	97	8.4	0.40	7.4	9.4	92	8.4	0.50	7.3	9.9	102	8.8*	0.32	8.0	9.8
Starch (Dry Basis %)	97	72.1	0.42	71.0	73.4	92	72.2	0.46	70.8	73.1	102	71.8*	0.41	70.5	72.4
Oil (Dry Basis %)	97	3.9	0.11	3.7	4.3	92	4.0*	0.12	3.6	4.2	102	3.9	0.14	3.5	4.4
Southern Rail						Southern Rail					Southern Rail				
Protein (Dry Basis %)	85	8.6	0.49	7.8	10.0	86	8.7	0.45	7.7	9.4	85	9.0*	0.43	8.2	10.0
Starch (Dry Basis %)	85	72.1	0.44	70.8	73.2	86	72.0	0.44	71.1	73.3	85	71.7*	0.46	70.7	72.8
Oil (Dry Basis %)	85	3.8	0.15	3.5	4.1	86	3.8	0.13	3.5	4.2	85	3.8	0.15	3.5	4.2

*Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

SUMMARY: CHEMICAL COMPOSITION

	Five-Year Average (MY20/21-MY24/25)			Ten-Year Average (MY15/16-MY24/25)			2025 Harvest					
	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.	No. of Samples ¹	Avg.	Std. Dev.	Min.	Max.	
U.S. Aggregate				U.S. Aggregate			U.S. Aggregate					
Protein (Dry Basis %)	2,190	8.6*	0.34	4,326	8.5*	0.32	621	8.4**	0.53	5.4	11.1	
Starch (Dry Basis %)	2,190	72.0*	0.40	4,326	72.3*	0.41	621	72.3**	0.55	70.2	74.1	
Oil (Dry Basis %)	2,190	3.9*	0.15	4,326	3.9*	0.15	621	3.8**	0.23	2.9	5.0	
Gulf				Gulf			Gulf					
Protein (Dry Basis %)	1,234	8.6*	0.29	2,577	8.5*	0.28	594	8.4**	0.54	5.4	11.1	
Starch (Dry Basis %)	1,234	72.0*	0.38	2,577	72.4*	0.38	594	72.3**	0.58	70.2	74.1	
Oil (Dry Basis %)	1,234	3.9	0.15	2,577	3.9*	0.15	594	3.8**	0.25	2.9	5.0	
Pacific Northwest				Pacific Northwest			Pacific Northwest					
Protein (Dry Basis %)	558	8.6*	0.40	1,041	8.6*	0.38	271	8.2**	0.51	5.4	11.1	
Starch (Dry Basis %)	558	71.8*	0.43	1,041	72.1	0.47	271	72.4**	0.47	70.2	73.8	
Oil (Dry Basis %)	558	3.9	0.14	1,041	4.0*	0.15	271	3.9	0.21	3.2	4.7	
Southern Rail				Southern Rail			Southern Rail					
Protein (Dry Basis %)	398	8.7	0.43	708	8.6	0.39	351	8.8	0.51	7.0	11.1	
Starch (Dry Basis %)	398	71.9*	0.43	708	72.2*	0.45	351	72.2**	0.54	70.2	74.1	
Oil (Dry Basis %)	398	3.8	0.16	708	3.9*	0.16	351	3.8**	0.22	2.9	4.6	





*Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

**Indicates current year's Export Cargo average was significantly different from this year's Harvest, based on a 2-tailed t-test at the 95% level of confidence.

¹Due to the ECA results being composite statistics, the sum of the sample numbers from the three ECAs is greater than the U.S. Aggregate.




**CHEMICAL COMPOSITION
AGGREGATE SIX-YEAR COMPARISON**



C. PHYSICAL FACTORS

Physical factors are other quality attributes that are neither grade factors nor chemical composition. Physical factors include stress cracks, kernel weight, kernel volume, true density, percent whole kernels and percent horneous (hard) endosperm. Tests for these physical factors provide additional information about the processing characteristics of corn for various uses and corn's storability and potential for breakage in handling. These quality attributes are influenced by the physical composition of the corn kernel, which is, in turn, affected by genetics and growing and handling conditions. Corn kernels are made up of four parts: the germ or embryo, the tip cap, the pericarp or outer covering and the endosperm. The endosperm represents about 82% of the kernel. It consists of soft (also referred to as floury or opaque) endosperm and of horneous (also called hard or vitreous) endosperm, as shown in the figure. The endosperm contains primarily starch and protein, the germ contains oil and some protein, and the pericarp and tip cap are mostly fiber.



Source: Adapted from Corn Refiners Association, 2011



SUMMARY: PHYSICAL FACTORS

- Average U.S. Aggregate stress cracks (15.0%) was lower than 2023/2024 (16.3%) but higher than 2024/2025 (11.6%), the 5YA (11.1%) and the 10YA (9.4%).
- Average U.S. Aggregate 100-kernel weight (36.83 g) was similar to 2024/2025 (36.95 g) and 2023/2024 (36.72 g) but higher than the 5YA (36.46 g) and the 10YA (36.00 g).
- Average 100-kernel weight for the Gulf ECA (38.18 g) was higher than the Southern Rail (35.89) and the Pacific Northwest (34.23 g) ECAs. Among ECAs, the Pacific Northwest has consistently had the lowest 100-kernel weight in the last five years, the 5YA and the 10YA.
- In 2025/2026, 84.1% of the samples had a 100-kernel weight of 34.00 g or higher, compared with 86.1% in 2024/2025 and 83.2% in 2023/2024.
- Average U.S. Aggregate kernel volume in 2025/2026 (0.29 cm³) was the same as 2024/2025, similar to 2023/2024 and 5YA (both 0.28 cm³) but statistically different (higher) than 10YA (0.28 cm³).
- Average kernel volume was lower for the Pacific Northwest ECA (0.27 cm³) than for the Gulf (0.29 cm³) and the Southern Rail ECAs (0.28 cm³) in 2025/2026. The Pacific Northwest ECA also had the lowest average kernel volume among ECAs for the five previous years, the 5YA and the 10YA.
- Average U.S. Aggregate kernel true density (1.289 g/cm³) was lower than 2024/2025 (1.293 g/cm³), similar to 2023/2024 (1.292 g/cm³) but higher than the 5YA and the 10YA (both 1.282 g/cm³).
- For the 2025/2026 export samples, 82.4% had kernel true densities equal to or above 1.275 g/cm³, compared to 93.3% in 2024/2025 and 84.2% in 2023/2024. The 2025/2026 samples contained fewer high-density kernels than in the previous two years.
- Previous to this year, the Pacific Northwest ECA has consistently had the lowest or tied for the lowest true densities and lowest test weights among ECAs for the past five years, the 5YA and the 10YA.
- The average percent of whole kernels at export (82.6%) was lower than 2024/2025 (89.1%), 2023/2024 (88.0%), the 5YA (86.3%) and the 10YA (85.6%).

- The percentage of 2025/2026 export samples with whole kernels greater than or equal to 85.0% was 36.5% as compared to 82.3% for 2024/2025 and 73.9% for 2023/2024.
- The Pacific Northwest ECA (79.7%) had the lowest average whole kernels compared to the Gulf (82.4%) and Southern Rail (86.4%) ECAs in 2025/2026. Among ECAs, the PNW also had the lowest whole kernels in the previous three years, the 5YA and the 10YA.
- Average U.S. Aggregate horneous endosperm (84%) was lower than 2024/2025 (87%) and 2023/2024 (86%), the same as the 5YA (84%) but higher than the 10YA (82%).
- Of the export samples, 97.8% in 2025/2026 had at least 80% horneous endosperm compared to 100.0% in 2024/2025 and 2023/2024.

STRESS CRACKS

Stress cracks are internal fissures in the horneous (hard) endosperm of a corn kernel. The pericarp (or outer covering) of a stress-cracked kernel is typically not damaged, so the kernel may appear unaffected at first glance, even if stress cracks are present.

The cause of stress cracks is pressure buildup due to moisture and temperature gradients within the kernel’s horneous endosperm. This can be likened to the internal cracks that appear when an ice cube is dropped into a lukewarm beverage. The internal stresses do not build up as much in the soft, floury endosperm as in the hard, horneous endosperm. Therefore, corn with a higher percentage of horneous endosperm is more susceptible to stress cracking than softer grain. A kernel may vary in severity of stress cracking and can have one, two or multiple stress cracks. The most common cause of stress cracks is high-temperature drying that rapidly removes moisture. The impact of high levels of stress cracks on various uses includes:

General: Increased susceptibility to breakage during handling. This may lead to processors needing to remove more broken corn during cleaning operations and a possible reduction in grade or value or both.


Wet Milling: Lower starch yields due to the increased difficulty in separating starch and protein. Stress cracks may also alter steeping requirements.

Dry Milling: The lower yield of large flaking grits (the prime product of many dry milling operations).

Alkaline Cooking: Non-uniform water absorption leading to overcooking or undercooking, which affects the process balance.

U.S. Aggregate Results Summary



STRESS CRACKS (%)



Growing conditions will affect crop maturity, the timing of harvest and the need for artificial drying, which will influence the degree of stress cracking found from region to region. For example, late maturity or late harvest caused by weather-related factors, such as rain-delayed planting or cool temperatures, may increase the need for artificial drying.

RESULTS: STRESS CRACKS

- Average U.S. Aggregate stress cracks (15.0%) was lower than 2023/2024 (16.3%) but higher than 2024/2025 (11.6%), the 5YA (11.1%) and the 10YA (9.4%).
- Average U.S. Aggregate stress cracks (15.0%) was higher than the 2025 harvest samples (9.5%).
- The standard deviation at export (8.4%) was similar to that at harvest (8.3%).
- Of the 2025/2026 export samples, 34.2% had 15% or higher stress cracks, compared with 29.2% in 2024/2025 and 50.8% in 2023/2024.
- The Pacific Northwest ECA (27.1%) had the highest average stress cracks followed by the Southern Rail ECA (15.6%) and the Gulf (10.1%)




100-KERNEL WEIGHT


Increased 100-kernel weight (reported in grams) indicates a larger kernel size. Kernel size affects drying rates. As kernel size increases, the volume-to-surface area ratio becomes higher, and as the ratio gets higher, drying becomes slower. In addition, large, uniform-sized kernels often enable higher flaking grit yields in dry milling.

RESULTS

- Average U.S. Aggregate 100-kernel weight (36.83 g) was similar to 2024/2025 (36.95 g) and 2023/2024 (36.72 g) but higher than the 5YA (36.46 g) and the 10YA (36.00 g).
- Average 100-kernel weight for export (36.83 g) was higher than at harvest (36.04 g). From 2011/2012 through 2024/2025, average 100-kernel weights ranged from 0.00 to 2.9 g greater at export than at harvest. Since 100-kernel weight is based on 100 fully intact kernels, any breakage or reduction in whole kernels occurring in transit may have self-selected out small kernels with low 100-kernel weights that might have been more prone to breakage.
- The export samples had a lower standard deviation (1.73 g) than the 2025 harvest samples (3.54 g). The 100-kernel weight standard deviation was also lower at export than at harvest for the last five years, the 5YA and the 10YA, indicating greater uniformity at export than at harvest.




- Average 100-kernel weight for the Gulf ECA (38.18 g) and the Southern Rail (35.89 g) were higher than the Pacific Northwest (34.23 g) ECA. Among ECAs, the Pacific Northwest consistently had the lowest 100-k weight in the last five years, the 5YA and the 10YA.
- In 2025/2026, 84.1% of the samples had a 100-kernel weight of 34.00 g or higher, compared with 86.1% in 2024/2025 and 83.2% in 2023/2024.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26



KERNEL VOLUME


Kernel volume, measured in cubic centimeters (cm³), is often indicative of growing conditions. If conditions are dry, kernels may be smaller than average. If a drought hits later in the season, kernels may have lower fill. Small or round kernels are more difficult to degerm. Additionally, small kernels may lead to increased cleanout losses for processors and higher yields of fiber.

RESULTS




- Average U.S. Aggregate kernel volume in 2025/2026 (0.29 cm³) was the same as 2024/2025, similar to 2023/2024 and 5YA (both 0.28 cm³) but statistically different (higher) than the 10YA (0.28 cm³).
- Kernel volume range (0.24 to 0.32 cm³) was similar to 2024/2025 (0.25 to 0.33 cm³), and 2023/2024 (0.21 to 0.32 cm³).
- The kernel volume standard deviation (0.01 cm³) was lower than 2023/2024 (0.02 cm³) but the same as 2024/2025, 5YA and the 10YA (all 0.01 cm³).
- Average U.S. Aggregate kernel volume at export (0.29 cm³) was higher than that for the 2025 harvest (0.27 cm³).

U.S. Aggregate Results Summary

KERNEL VOLUME (cm³)



- Average kernel volume was lower for the Pacific Northwest ECA (0.27 cm³) than for the Gulf (0.29 cm³) and the Southern Rail ECA (0.28 cm³) in 2025/2026. The Pacific Northwest ECA also had the lowest average kernel volume among ECAs for the five previous years, the 5YA and the 10YA.
- Of the 2025/2026 export samples, 43.9% had kernel volumes equal to or higher than 0.29 cm³, compared to 46.1% in 2024/2025 and 45.7% in 2023/2024.
- There is a positive relationship between kernel volume and 100-kernel weight in the 2025/2026 export samples, as shown in the adjacent figure (the correlation coefficient is 0.98). This indicates the larger the kernel size, the higher the 100-kernel weight.




KERNEL TRUE DENSITY


Kernel true density is calculated as the weight of a 100-kernel sample divided by the volume or displacement of those 100 kernels and is reported as grams per cubic centimeter (g/cm³). True density is a relative indicator of kernel hardness, which is useful for alkaline processors and dry millers. True density may be affected by the genetics of the corn hybrid and the growing environment. Corn with a higher density is typically less susceptible to breakage in handling than lower density corn but is also more at risk of developing stress cracks if high-temperature drying is employed. True densities above 1.30 g/cm³ indicate very hard corn, which is typically desirable for dry milling and alkaline processing. True densities near the 1.275 g/cm³ level and below tend to be softer but process well for wet milling and feed use.

RESULTS

- Average U.S. Aggregate kernel true density for the 2025/2026 export samples (1.289 g/cm³) was lower than 2024/2025 (1.293 g/cm³), similar to 2023/2024 (1.292 g/cm³) but higher than the 5YA and the 10YA (both 1.282 g/cm³).
- Average kernel true density for the 2025/2026 export samples (1.289 g/cm³) was higher than for the 2025 harvest samples (1.258 g/cm³).




- The 2025/2026 export samples had a lower range (1.233 to 1.328 g/cm³) than the 2025 harvest samples (1.194 to 1.323 g/cm³). Similarly, the standard deviation of this year's export samples (0.014 g/cm³) was much lower than the standard deviation at harvest (0.025 g/cm³).
- For the 2025/2026 export samples, 82.4% had kernel true densities equal to or above 1.275 g/cm³, compared to 93.3% in 2024/2025 and 84.2% in 2023/2024. This indicates the samples contained fewer kernels with high density in 2025/2026 than in the previous two years.
- Average kernel true densities for ECAs were 1.298 g/cm³ for the Gulf, 1.279 g/cm³ for the Pacific Northwest and 1.277 g/cm³ for the Southern Rail. Previous to this year, the Pacific Northwest has had the lowest or tied for the lowest true densities and lowest test weights among ECAs for the previous four years, the 5YA and the 10YA.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26



WHOLE KERNELS

The whole kernels test measures the percent of fully intact kernels in the sample with no pericarp damage or kernel pieces chipped away. The exterior integrity of the corn kernel is very important for two key reasons. First, it affects water absorption for alkaline cooking and steeping operations. Kernel nicks or pericarp cracks allow water to enter the kernel faster than intact or whole kernels. Too much water uptake during cooking can result in loss of solubles, non-uniform cooking, expensive shutdown time, or products that do not meet specifications. Some companies pay contracted premiums for corn delivered above a specified level of whole kernels.


Second, intact whole kernels are less susceptible to storage molds and breakage in handling. While hard endosperm corn lends itself to the preservation of more whole kernels than soft endosperm corn, the primary factor in delivering whole kernels is harvesting and handling. This begins with proper combine adjustment, followed by minimizing the severity of kernel impacts due to conveyors and the number of handlings required from the farm field to the end-user. Each subsequent handling will generate additional breakage. Actual amounts of breakage increase exponentially as moisture decreases, drop heights increase or a kernel's velocity at impact increases.¹ In addition, harvesting at the higher moisture content (e.g., greater than 25%) will usually lead to soft pericarps and more pericarp damage to corn than when harvesting at lower moisture levels.

RESULTS

- Average U.S. Aggregate whole kernels (82.6%) was lower than 2024/2025 (89.1%), 2023/2024 (88.0%), the 5YA (86.3%) and the 10YA (85.6%).


WHOLE KERNELS (%)

U.S. Aggregate Results Summary




¹Foster, G.H. and L.E. Holman. 1973. Grain Breakage Caused by Commercial Handling Methods. Marketing Research Report No. 968. ARS, USDA, Washington, D.C.

- The average percentage of whole kernels at export in 2025/2026 (82.6%) was lower than at harvest (90.6%). Average whole kernels for the export 5YA (86.3%) was also lower than for the harvest 5YA (92.3%).
- Over the past five years and for the 5YA and 10YA, the percentages of whole kernels at export have been 3.1 to 9.3 percentage points lower than at harvest. This reduction in whole kernels is likely caused by the additional handling during transport to export loading locations.
- The 2025/2026 export samples had a range of 53.0 to 96.2% whole kernels and a standard deviation of 5.9%.
- The Pacific Northwest ECA (79.7%) had the lowest average whole kernels compared to the Gulf (82.4%) and Southern Rail (86.4%) ECAs in 2025/2026. Among ECAs, the PNW also had the lowest whole kernels in the previous three years, the 5YA and the 10YA.
- The percentage of export samples with whole kernels greater than or equal to 85% was 36.5% for 2025/2026, 82.3% for 2024/2025 and 73.9% for 2023/2024.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26



HORNEOUS (HARD) ENDOSPERM

The horneous (hard) endosperm test measures the percent of horneous or hard endosperm out of the total endosperm in a kernel, with a potential value from 70 to 100%. The greater the amount of horneous endosperm relative to soft endosperm, the harder the corn kernel is said to be. The degree of hardness is important, depending on the type of processing. A hard kernel is needed to produce high yields of large-flaking grits in dry milling. Hard to medium hardness is desired for alkaline cooking. Medium to soft hardness is used for wet milling and livestock feeding.


Hardness has been correlated to breakage susceptibility, feed utilization/efficiency and starch digestibility. As a test of overall hardness, there is no good or bad value for horneous endosperm. There is only a preference by different end-users for particular ranges. Many dry millers and alkaline cookers would like greater than 85% horneous endosperm, while wet millers and feeders would typically like values between 70% and 85%. However, there are certainly exceptions in user preference.

RESULTS


- Average U.S. Aggregate horneous endosperm (84%) was lower than 2024/2025 (87%) and 2023/2024 (86%), the same as 5YA (84%) but higher than the 10YA (82%).
- Horneous endosperm at export (84%) was higher than that at harvest (83%).

HORNEOUS ENDOSPERM (%)

U.S. Aggregate Results Summary




- The 2025/2026 export samples had more uniform percentages of horneous endosperm than the 2025 harvest samples, as indicated by the lower standard deviation at export (2%) compared to that at harvest (3%).
- The export samples also had a range (79 to 90%) which was less than the harvest samples (74 to 92%). This same pattern of increased uniformity for the export samples compared to the harvest samples also occurred in the three previous years, the 5YA and the 10YA.
- Average horneous endosperm was 84% for the Gulf and Southern Rail ECAs and 83% for the Pacific Northwest ECA.
- Of the 2025/2026 export samples, 97.8% had 80% or higher horneous endosperm, compared to 100.0% in 2024/2025 and in 2023/2024.



Percent of Samples by Marketing Year

■ MY23/24 ■ MY24/25 ■ MY25/26





SUMMARY: PHYSICAL FACTORS

2025/2026 Export Cargo						2024/2025 Export Cargo					2023/2024 Export Cargo				
	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.	No. of Samples	Avg.	Std. Dev.	Min.	Max.
U.S. Aggregate						U.S. Aggregate					U.S. Aggregate				
Stress Cracks (%)	430	15.0	8.4	1	66	425	11.6*	7.0	0	51	433	16.3*	8.5	0	47
100-Kernel Weight (g)	182	36.83	1.73	30.97	41.40	180	36.95	1.38	31.38	42.24	184	36.72	2.44	26.24	41.83
Kernel Volume (cm ³)	182	0.29	0.01	0.24	0.32	180	0.29	0.01	0.25	0.33	184	0.28	0.02	0.21	0.32
True Density (g/cm ³)	182	1.289	0.014	1.233	1.328	180	1.293*	0.009	1.248	1.320	184	1.292	0.011	1.244	1.319
Whole Kernels (%)	430	82.6	5.9	53.0	96.2	425	89.1*	4.4	67.8	97.8	433	88.0*	4.4	69.8	99.6
Horneous Endosperm (%)	182	84	2	79	90	180	87*	2	81	92	184	86*	1	83	90
Gulf						Gulf					Gulf				
Stress Cracks (%)	248	10.1	4.5	1	24	247	11.7*	6.0	1	37	246	17.5*	9.1	4	47
100-Kernel Weight (g)	104	38.18	1.60	34.35	41.40	108	37.89	1.31	34.14	42.24	108	37.99	2.16	31.77	41.83
Kernel Volume (cm ³)	104	0.29	0.01	0.27	0.32	108	0.29	0.01	0.26	0.33	108	0.29	0.02	0.25	0.32
True Density (g/cm ³)	104	1.298	0.010	1.275	1.328	108	1.300	0.009	1.272	1.320	108	1.300	0.008	1.277	1.319
Whole Kernels (%)	248	82.4	4.8	67.8	94.8	247	88.7*	4.5	67.8	97.8	246	88.0*	4.2	73.6	99.6
Horneous Endosperm (%)	104	84	2	81	89	108	87*	2	81	92	108	86*	1	83	90
Pacific Northwest						Pacific Northwest					Pacific Northwest				
Stress Cracks (%)	97	27.1	13.3	7	66	92	14.2*	10.5	1	42	102	17.5*	7.8	3	40
100-Kernel Weight (g)	42	34.23	1.58	30.97	38.40	36	33.55*	1.04	31.38	35.05	37	33.12*	2.89	26.24	37.13
Kernel Volume (cm ³)	42	0.27	0.01	0.24	0.31	36	0.26*	0.01	0.25	0.28	37	0.26*	0.02	0.21	0.29
True Density (g/cm ³)	42	1.279	0.014	1.245	1.308	36	1.284	0.008	1.266	1.311	37	1.279	0.015	1.244	1.305
Whole Kernels (%)	97	79.7	7.6	56.8	94.6	92	88.4*	4.6	78.6	96.8	102	84.3*	5.5	69.8	96.2
Horneous Endosperm (%)	43	83	2	79	86	36	86*	2	81	88	37	86*	0	85	87
Southern Rail						Southern Rail					Southern Rail				
Stress Cracks (%) ¹	85	15.6	14.5	1	62	86	8.8*	6.7	0	51	85	11.5*	7.1	0	39
100-Kernel Weight (g)	36	35.89	2.25	31.40	40.41	36	37.50*	1.92	32.14	40.78	39	36.70	2.83	30.75	40.54
Kernel Volume (cm ³)	36	0.28	0.02	0.24	0.31	36	0.29*	0.01	0.25	0.32	39	0.29	0.02	0.25	0.31
True Density (g/cm ³)	36	1.277	0.023	1.233	1.309	36	1.284	0.012	1.248	1.305	39	1.282	0.018	1.244	1.309
Whole Kernels (%)	85	86.4	7.0	53.0	96.2	86	90.9*	3.9	79.2	97.6	85	91.8*	3.6	83.0	98.8
Horneous Endosperm (%)	35	84	2	79	90	36	86*	2	81	90	39	86*	1	84	88

^{*}Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

¹The relative margin of error for predicting the population average exceeded 10.0%.

SUMMARY: PHYSICAL FACTORS

	Five-Year Average (MY20/21-MY24/25)			Ten-Year Average (MY15/16-MY24/25)			2025 Harvest				
	No. of Samples	Avg.	Std. Dev.	No. of Samples	Avg.	Std. Dev.	No. of Samples ¹	Avg.	Std. Dev.	Min.	Max.
U.S. Aggregate											
Stress Cracks (%)	2,187	11.1*	6.5	4,323	9.4*	5.8	621	9.5**	8.3	0	96
100-Kernel Weight (g)	924	36.46*	1.75	3,060	36.00*	1.65	182	34.04**	3.54	25.50	41.90
Kernel Volume (cm ³)	924	0.28	0.01	3,060	0.28*	0.01	182	0.27**	0.03	0.21	0.33
True Density (g/cm ³)	924	1.282*	0.012	3,060	1.282*	0.012	182	1.258**	0.025	1.194	1.323
Whole Kernels (%)	2,165	86.3*	4.5	4,301	85.6*	4.8	621	90.6**	4.9	66.4	99.4
Horneous Endosperm (%)	928	84	2	2,812	82*	2	182	83**	3	74	92
Gulf											
Stress Cracks (%)	1,225	11.3*	6.2	2,568	9.1*	5.4	594	9.9	8.9	0	96
100-Kernel Weight (g)	519	37.85	1.62	1,862	37.19*	1.56	172	34.70**	3.55	25.50	41.90
Kernel Volume (cm ³)	519	0.29	0.01	1,862	0.29*	0.01	172	0.28**	0.03	0.21	0.33
True Density (g/cm ³)	519	1.290*	0.010	1,862	1.288*	0.010	172	1.260**	0.025	1.194	1.323
Whole Kernels (%)	1,226	86.3*	4.3	2,569	86.1*	4.5	594	90.5**	5.1	66.4	99.4
Horneous Endosperm (%)	523	84	2	1,726	83*	2	172	83**	3	74	92
Pacific Northwest											
Stress Cracks (%)	558	12.8*	7.4	1,041	12.0*	6.7	271	10.7**	9.8	1	96
100-Kernel Weight (g)	226	32.95*	1.76	709	32.72*	1.68	80	32.51**	3.18	26.40	41.00
Kernel Volume (cm ³)	226	0.26*	0.01	709	0.26*	0.01	80	0.26	0.02	0.21	0.32
True Density (g/cm ³)	226	1.268*	0.013	709	1.271*	0.014	80	1.244**	0.026	1.194	1.302
Whole Kernels (%)	556	84.2*	5.3	1,039	82.7*	5.7	271	90.6**	4.7	73.2	98.8
Horneous Endosperm (%)	226	83*	1	639	82*	2	80	82	4	74	92
Southern Rail											
Stress Cracks (%)	404	8.7*	6.3	714	7.1*	6.0	351	6.7**	4.7	0	59
100-Kernel Weight (g)	179	36.60	2.14	489	36.33	1.90	99	33.95**	3.43	25.50	41.90
Kernel Volume (cm ³)	179	0.29*	0.02	489	0.28	0.01	99	0.27**	0.03	0.21	0.33
True Density (g/cm ³)	179	1.276	0.016	489	1.278	0.014	99	1.268	0.021	1.194	1.323
Whole Kernels (%)	383	89.1*	4.3	693	88.1*	4.5	351	91.1**	4.7	70.0	99.4
Horneous Endosperm (%)	179	84	2	447	82*	2	99	85	3	74	91





*Indicates average was significantly different from current year's Export Cargo, based on a 2-tailed t-test at the 95% level of confidence.

**Indicates current year's Export Cargo average was significantly different from this year's Harvest, based on a 2-tailed t-test at the 95% level of confidence.




¹Due to the ECA results being composite statistics, the sum of the sample numbers from the three ECAs is greater than the U.S. Aggregate.



**PHYSICAL FACTORS
AGGREGATE SIX-YEAR COMPARISON**



PHYSICAL FACTORS AGGREGATE SIX-YEAR COMPARISON



D. MYCOTOXINS

Mycotoxins are toxic compounds produced by fungi that occur naturally in grains. When consumed at elevated levels, mycotoxins may cause sickness in humans and animals. Several mycotoxins have been found in corn grain. Aflatoxin, DON and fumonisin are considered three of the important mycotoxins.

A subset of the export samples has been tested for aflatoxin and DON in all fifteen years of *Export Cargo Reports*. Beginning with the *2019/2020 Export Cargo Report*, fumonisins were added to the list of mycotoxins tested. In addition to testing survey samples for aflatoxin, DON and fumonisin, the *2021/2022 Export Cargo Report* also began testing export samples for ochratoxin A, T-2 and zearalenone. The testing for these three additional mycotoxins is intended to complement the information provided by the aflatoxin, DON and fumonisin test results and is continued in this *2025/2026 Export Cargo Report*.

Depending on the year, environmental conditions under which the corn is produced and stored may or may not be conducive to developing a particular mycotoxin to levels that impact the corn's use for human and livestock consumption. Humans and livestock are sensitive to mycotoxins at varying levels. As a result, the FDA has issued action levels for aflatoxin and advisory levels for DON and fumonisin by intended use.

Action levels specify limits of contamination above which the agency is prepared to take regulatory action. Action levels signal that the FDA believes it has data to support regulatory or court action if a toxin or contaminant is present at levels exceeding the action level if the agency chooses to do so. If imports or domestic feed supplements are analyzed in accordance with valid methods and found to exceed action levels, they are considered adulterated and may be seized and removed from interstate commerce by the FDA.

Advisory levels guide the industry concerning levels of a substance present in food or feed believed by the agency to provide an adequate margin of safety to protect human and animal health. While the FDA reserves the right to take regulatory enforcement action, enforcement is not the fundamental purpose of an advisory level. A source of additional information is the National Grain and Feed Association (NGFA) guidance document titled "FDA Mycotoxin Regulatory Guidance," found at the following link: https://drive.google.com/file/d/1tqeS5_eOtsRmxZ5RrTnYu7NClr896KGX/view.

The table below displays the mycotoxin test kits utilized for each of the six mycotoxins tested in the *2025/2026 Export Cargo Report* along with the minimum reported level for each mycotoxin and each test kit’s limit of detection. The limit of detection is a threshold indicating whether or not a detectable level of the mycotoxin appeared in the sample. Additional details on the testing methodology employed in this study for mycotoxins are in the “Testing Analysis Methods” section.

Mycotoxin	Export Cargo Report Minimum Reported Levels	Mycotoxin Test Kit (EnviroLogix)	Test Kit Limit of Detection
Aflatoxin	5.0 parts per billion	AQ 309 BG	2.7 parts per billion
DON	1.5 parts per million	AQ 304 BG	0.1 parts per million
Fumonisin	5.0 parts per million	AQ 411 BG	0.1 parts per million
Ochratoxin A	5.0 parts per billion	AQ 113 BG	1.5 parts per billion
T-2	1.5 parts per million	AQ 314 BG	50 parts per billion
Zearalenone	1.5 parts per million	AQ 412 BG	50 parts per billion

Source: www.envirologix.com

AFLATOXIN

The most important type of mycotoxin associated with corn grain is aflatoxin. There are several types of aflatoxin produced by different species of *Aspergillus*, with the most prominent species being *A. flavus*. The growth of the fungus and aflatoxin contamination of grain can occur in the field prior to harvest or in storage. However, contamination prior to harvest is considered to cause most of the problems associated with aflatoxin. *A. flavus* grows well in hot, dry environmental conditions or where drought occurs over an extended period. It can be a serious problem in the southern United States, where hot and dry conditions are more common. The fungus usually attacks only a few kernels on the ear and often penetrates kernels through wounds produced by insects. Under drought conditions, it also grows down silks into individual kernels.

There are four types of aflatoxin naturally found in foods – aflatoxins B1, B2, G1 and G2. These four aflatoxins are commonly referred to as “aflatoxin” or “total aflatoxin.” Aflatoxin B1 is the most commonly found aflatoxin in food and feed and is also the most toxic. Research has shown that B1 is a potent, naturally-occurring carcinogen in animals, with a strong link to human cancer incidence. Additionally, dairy cattle will metabolize aflatoxin to a different form of aflatoxin called aflatoxin M1, which may accumulate in milk.

Aflatoxin expresses toxicity in humans and animals primarily by attacking the liver. The toxicity can occur from short-term consumption of very high doses of aflatoxin-contaminated grain or long-term ingestion of low levels of aflatoxin, possibly resulting in death in poultry, the most sensitive of the animal species. Livestock may experience reduced feed efficiency or reproduction, and both human and animal immune systems may be suppressed as a result of ingesting aflatoxin.

The FDA has established action levels in parts per billion (ppb) for aflatoxin M1 in milk intended for human consumption and aflatoxin in human food, grain and livestock feed (see table below).

Aflatoxin Action Level	Criteria
20.0 parts per billion	Dairy animals, pets of all ages, immature animals (including immature poultry) and when the animal’s destination is not known
100.0 parts per billion	Breeding beef cattle, breeding swine and mature poultry
200.0 parts per billion	Finishing swine of 100 pounds or greater
300.0 parts per billion	Finishing (i.e., feedlot) beef cattle

Source: www.ngfa.org

For additional information, see the National Grain and Feed Association’s guidance document titled “FDA Mycotoxin Regulatory Guidance” found at https://drive.google.com/file/d/1tqeS5_eOtsRmxZ5RrTnYu7NC1r896KGX/view.


The FDA has established additional policies and legal provisions concerning the blending of corn with levels of aflatoxin exceeding these threshold levels. In general, the FDA currently does not permit the blending of corn blended to reduce the aflatoxin content to be sold in general commerce.

Corn exported from the United States must be tested for aflatoxin according to federal law. Unless the contract exempts this requirement, testing must be conducted by FGIS. Corn above the FDA action level of 20.0 ppb cannot be exported unless other strict conditions are met. This results in relatively low levels of aflatoxin in exported grain.

RESULTS

A total of 182 export samples was tested by USDA FGIS for aflatoxin for the *2025/2026 Export Cargo Report*. A threshold established by USDA FGIS as the “Lower Conformance Level” was used to determine whether or not a detectable level of aflatoxin appeared in the sample. The Lower Conformance Level for aflatoxin of the FGIS-approved analytical kits used for this 2025/2026 report was 5.0 ppb. Results of the 2025/2026 survey testing are as follows:

- Of the 182 samples tested, 175 samples (96.2%) had no detectable levels of aflatoxin (below the FGIS LCL of 5.0 ppb). This 96.2% is similar to 2024/2025 (96.1%) and higher than 2023/2024 (91.2%).
- Two (2) samples (1.1%) were found with aflatoxin levels greater than or equal to 5.0 ppb, but less than 10.0 ppb of the 182 samples tested in 2025/2026. This percentage is lower than 2024/2025 (3.3%) and 2023/2024 (7.2%).
- Two (2) samples (1.1%) of the 182 samples tested in 2025/2026 had aflatoxin levels greater than or equal to 10.0 ppb, but below or equal to the FDA action level of 20.0 ppb. This 1.1% is greater than 2024/2025 (0.6%) and less than 2023/2024 (1.7%).
- Three (3) samples (1.6%) tested in 2025/2026 were above the FDA action level of 20.0 ppb, which is greater than 2024/2025 (0.0%) and 2023/2024 (0.0%).



As in the previous two years, aflatoxin contamination in 2025/2026 may be indicative of weather conditions during the 2025 growing season that were not conducive to aflatoxin formation.

DEOXYNIVALENOL (DON OR VOMITOXIN)

DON is another mycotoxin of concern to some importers of corn grain. It is produced by a certain species of *Fusarium*, the most important of which is *Fusarium graminearum* (*Gibberellazeae*), which also causes Gibberella ear rot (or red ear rot). *Gibberellazeae* can develop when cool or moderate and wet weather occurs at flowering. The fungus grows down the silks into the ear. In addition to producing DON, it produces conspicuous red discoloration of kernels on the ear. The fungus can also continue to grow and rot ears when corn is left standing in the field. Therefore, mycotoxin contamination of corn caused by *Gibberellazeae* is often associated with excessive postponement of harvest and/or storage of high-moisture corn.

DON is mostly a concern with monogastric animals, where it may cause irritation of the mouth and throat. As a result, animals may eventually refuse to eat the DON-contaminated corn and may have low weight gain, diarrhea, lethargy and intestinal hemorrhaging. In addition, it may cause suppression of the immune system, resulting in susceptibility to a number of infectious diseases.

The FDA has issued advisory levels for DON in parts per million (ppm). For products containing corn, the advisory levels are shown below.

DON Advisory Level	Criteria
5.0 parts per million	Swine, not to exceed 20% of their diet
5.0 parts per million	All other animals not otherwise listed, not to exceed 40% of their diet
10.0 parts per million	Chickens, not to exceed 50% of their diet
10.0 parts per million	Ruminating beef and dairy cattle older than four months

Source: www.ngfa.org

For additional information, see the National Grain and Feed Association's guidance document titled "FDA Mycotoxin Regulatory Guidance" found at https://drive.google.com/file/d/1tqeS5_eOtsRmxZ5RrTnYu7NCIr896KGX/view.


FGIS is not required to test for DON on corn bound for export markets but will perform either a qualitative or quantitative test for DON at the buyer's request.

RESULTS

A total of 182 export samples was tested for DON for the *2025/2026 Export Cargo Report*. Results of the testing are shown below:

- One hundred seventy (170) of the 182 samples (93.4%) tested less than 1.5 ppm. This 93.4% is lower than 2024/2025 (96.7%) but the same as 2023/2024.
- Eleven (11) samples of the 182 samples (6.0%) tested in 2025/2026 had DON levels greater than or equal to 1.5 ppm but less than 5.0 ppm. This 6.0% is higher than 2024/2025 (3.3%) but similar to 2023/2024 (6.6%).
- One (1) of the 182 samples tested in 2025/2026 (0.5%) was above the FDA advisory level of 5.0 ppm, which is higher than 2024/2025 and 2023/2024 (both 0.0%).

While one sample was above the FDA advisory level of 5.0 ppm in 2025/2026, these results were indicative of the excellent growing season that was not conducive to DON contamination.



FUMONISIN

Fumonisin is a naturally occurring mycotoxin found mostly in cereal grains, mainly corn. Fumonisin is a more recent discovery compared to aflatoxin and DON. Fumonisin is produced by several fungi of the *Fusarium* genus. The fumonisin family consists of fumonisin B1, fumonisin B2 and fumonisin B3. Fumonisin B1 is the most abundant, accounting for about 70 to 80% of the sum of the three fumonisins. The main concern with fumonisin is feed contamination that can have detrimental effects, particularly on horses and pigs. Fungal and fumonisin formation occurs mainly before harvest. Insects play an important role in fumonisin contamination since they act as a wounding agent. Temperature and rainfall conditions are related to fungal growth and fumonisin contamination. In general, fumonisin contamination is related to plant stress, insect damage, drought and soil moisture. In 2001, FDA issued guidance levels for the sum of the three fumonisins in corn-based foods and feed to reduce human and animal exposure. FDA advisory levels are shown below.

Fumonisin Advisory Level	Criteria
5.0 parts per million	Equids (i.e., horses) and rabbits, not to exceed 20% of diet
20.0 parts per million	Swine and catfish, not to exceed 50% of diet
30.0 parts per million	Breeding ruminants, breeding poultry and breeding mink, not to exceed 50% of diet
60.0 parts per million	Ruminants older than three months raised for slaughter and mink raised for pelt production, not to exceed 50% of diet
100.0 parts per million	Poultry raised for slaughter, not to exceed 50% of diet
10.0 parts per million	All other animals not otherwise listed, not to exceed 50% of their diet

Source: www.ngfa.org


For additional information, see the National Grain and Feed Association's guidance document titled "FDA Mycotoxin Regulatory Guidance" found at https://drive.google.com/file/d/1tqeS5_eOtsRmxZ5RrTnYu7NC1r896KGX/view.

RESULTS

A total of 182 samples was analyzed for fumonisin in the 2025/2026 report. This is the seventh year that survey samples have been tested for fumonisin. Results of the 2025/2026 survey are as follows:

- One hundred seventy-six (176) of the 182 samples (96.7%) tested below 5.0 ppm, the lowest advisory level for animals (equids and rabbits), which is similar to 2024/2025 (97.8%) and 2023/2024 (97.2%).
- One (1) of the 182 samples (0.5%) tested greater than or equal to 5.0 ppm, but less than 10.0 ppm, which is lower than 2024/2025 and 2023/2024 (both 2.2%).
- Four (4) of the 182 samples (2.2%) tested greater than or equal to 10.0 ppm, but less than 30.0 ppm, which is greater than 2024/2025 (0.0%) and 2023/2024 (0.6%).
- One (1) of the 182 samples (0.5%) tested greater than 30.0 ppm, which is the advisory level for breeding ruminants, poultry and mink, and is greater than 2024/2025 and 2023/2024 (both 0.0%).

All but 1 sample (0.5%) in 2025/2026 survey results tested below the 30.0 ppm level which is the advisory level for breeding ruminants, poultry and mink. This is likely indicative of the weather conditions during the 2025 growing season, which were not conducive for mold growth and fumonisin formation.



OCHRATOXIN A


Ochratoxins are considered a hazardous mycotoxin produced by a number of fungal species such as *Penicillium verrucosum* and *Aspergillus ochraceus* that can colonize grains, cereals and a range of other food products. Of these products, grains and cereals are considered to represent 50–80% of the intake of ochratoxins. The fungi can produce ochratoxins A, B and C, but ochratoxin A is produced in the greatest quantity. While ochratoxin A can occur all along the production chain from the field to storage, it is primarily considered a storage problem. Grains stored under high moisture/humidity (>14%) at warm temperatures (>20°C) and/or inadequately dried have the potential to become contaminated with the fungi and produce ochratoxins. Also, damage to the grain by mechanical means, physical means or insects can provide a portal of entry for the fungus. Initial growth of fungi in grains can form sufficient moisture from metabolism to allow for further growth and mycotoxin formation. Because grains and cereal products represent a large portion of the human diet, several countries have established maximum levels for ochratoxin A in unprocessed cereals. The European Commission established a maximum level for ochratoxin A in raw cereals at 5.0 parts per billion. The FDA has issued no advisory levels for ochratoxin A.

This is the fifth year that export samples were tested for ochratoxin A. As with the other mycotoxins, 182 samples were tested to assess the impact of this year’s conditions under which the corn was produced and stored on ochratoxin A. The testing methodology employed is described in the “Testing Analysis Methods” section.

RESULTS

Results of the 182 samples analyzed for ochratoxin A in the 2025/2026 survey are as follows:

- One hundred eighty-two (182) or 100% of the samples tested were below 5.0 ppb, the European Commission’s established maximum level for ochratoxin A. This percentage was higher than 2024/2025 (98.3%) and 2023/2024 (97.8%).
- None (0) or 0.0% of the 182 samples tested were greater than or equal to 5.0 ppb, but not greater than 10.0 ppb. This percentage was lower than 2024/2025 and 2023/2024 (both 1.7%).
- None (0) or 0.0% of the 182 samples tested were greater than 10.0 ppb. This percentage was the same as 2024/2025 (0.0%) and lower than 2023/2024 (0.6%).
- These results were indicative of the excellent growing season that was not conducive to OTA contamination.



T-2

T-2 is one of several mycotoxins (including Deoxynivalenol or DON) belonging to a group of mycotoxins called trichothecenes. T-2 toxin is produced in growing cereal grain crops by various *Fusarium* species of fungi. The fungi can grow over a wide range of temperatures (-2 to 35°C) and only at a water activity above 0.88. As a result, T-2 is not normally found in grain at harvest but in grain that has suffered water damage when left in the field after harvest (especially over winter). However, T-2 can occur in storage if the grain has suffered water damage in storage. The FDA has issued no advisory levels for T-2 toxin.


This is the fifth year that export samples were tested for T-2. A total of 182 samples were tested to assess the impact of this year’s conditions under which the corn was produced and stored on T-2. The testing methodology employed is described in the “Testing Analysis Methods” section.

RESULTS

Results of the 182 samples analyzed for T-2 in the 2025/2026 survey are as follows:

- One hundred eighty-two (182) or 100.0% of the samples tested were below 1.5 ppm. This 2025/2026 percentage (100.0%) was the same as 2024/2025 (100%) and 2023/2024 (100%).
- None (0) or 0.0% of the 182 samples tested greater than or equal to 1.5 ppm, but not greater than 5.0 ppm. This 2025/2026 percentage (0.0%) was the same as 2024/2025 and 2023/2024.
- None (0) or 0.0% of the 182 samples test greater than 5.0 ppm. This 2025/2026 percentage (0.0%) was the same as 2024/2025 and 2023/2024.

These results were indicative of the excellent growing season that was not conducive to T-2 contamination.





ZEARALENONE


Zearalenone is a mycotoxin that is very similar to DON in most aspects, with a few exceptions. Both are produced by *Fusarium* species of fungi. As a result, it would not be uncommon to find both mycotoxins in grain and grain products at the same time. The growing conditions for zearalenone production are very comparable to DON, with the optimal temperatures ranging from 65 to 85°F. A drop in temperature during growth also stimulates the production of toxins by the fungi. A moisture content of 20% or greater is required by the fungi to produce zearalenone, which is also similar to that needed to produce DON. But if the moisture content during growth drops below 15%, the production of toxins is halted. This is one of the reasons it is recommended that corn for storage should be dried to moisture levels less than 15%. Levels of as little as 0.1 ppm to 5.0 ppm have been shown to cause reproductive problems in swine, so great care should be used when feeding possibly contaminated grain to pigs. The FDA has issued no advisory levels for zearalenone but recommends only that the levels of concern for DON be observed.

As with the other mycotoxins, 182 samples were tested to assess the impact of this year’s conditions under which the corn was produced and stored on zearalenone. The testing methodology employed is described in the “Testing Analysis Methods” section.

RESULTS

This is the fifth year that survey samples were tested for zearalenone. Results of the 182 samples analyzed in the 2025/2026 survey are as follows:

- One hundred eighty-two (182) or 100.0% of the samples tested were below 1.5 ppm. The 2025/2026 percentage (100.0%) was the same as 2024/2025 and 2023/2024.
- None (0) or 0.0% of the 182 samples tested were greater than or equal to 1.5 ppm, but not greater than 5.0 ppm. The 2025/2026 percentage (0.0%) was the same as 2024/2025 and 2023/2024.
- None (0) or 0.0% of the 182 samples tested were greater than 5.0 ppm. The 2025/2026 percentage (0.0%) was the same as 2024/2025 and 2023/2024.



This *2025/2026 Export Cargo Report* provides advanced information about corn quality by evaluating and reporting quality attributes when the corn is ready to be loaded onto the ocean-going vessel or railcar for export. Corn quality includes a range of attributes that can be categorized as:


- Intrinsic quality characteristics – Protein, oil and starch concentrations as well as kernel hardness and density are intrinsic quality characteristics, meaning they are contained within and are of critical importance to the end-user. Since they are non-visual, they can only be determined by analytical tests.
- Physical quality characteristics – These attributes are associated with the outward visible appearance of the kernel or measurement of the kernel characteristics. Characteristics include kernel size, shape and color; test weight; total damaged and heat-damaged kernels; broken kernels; and stress cracks. Some of these characteristics are measured when corn receives an official USDA grade.
- Sanitary quality characteristics – These characteristics indicate the cleanliness of the grain. Attributes include foreign material, odor, dust, rodent excreta, insects, residues, fungal infection and non-millable materials.

The intrinsic quality characteristics are impacted significantly by genetics and growing season conditions and typically do not change at the aggregate level as corn moves through the marketing system. If the measured values of the intrinsic characteristics differ between harvest and export at the aggregate level, the differences can be due partially to normal random variation in sampling. On the other hand, the physical and sanitary characteristics can change as corn moves through the marketing channel. The parties involved in corn marketing and distribution use operating practices (such as cleaning, drying and conditioning) at each step in the channel to increase uniformity, prevent or minimize the loss of physical and sanitary quality and meet contract specifications.

The *Harvest Report* assesses the quality of the recently harvested corn crop as it enters the marketing system. The *Export Cargo Report* provides information on the impact of subsequent practices, including cleaning, drying, handling, blending, storing and transporting the crop up to the point where it is being loaded for export. To provide the backdrop for this assessment, the following sections describe the marketing channel from farm to export, the practices applied to corn as it moves through the marketing channel and the implications of these practices on corn quality. Lastly, the inspection and grading services provided by USDA FGIS or an official service provider are reviewed.

A. U.S. CORN EXPORT FLOW

As corn is harvested, farmers transport grain to on-farm storage, end-users or commercial grain facilities. While some producers feed their corn to their own livestock, most of the corn moves to other end-users (feed mills or processors) or commercial grain-handling facilities, such as local grain elevators, inland subterminals, river elevators and port elevators. Local grain elevators typically receive most of their grain directly from farmers. Inland subterminals or river elevators collect grain in quantities suitable for loading on unit trains and barges for further transport. These elevators may receive more than half of their corn from other elevators and are often located where the transport of bulk grain can be easily accommodated by unit trains or barges. Local grain, inland subterminals and river elevators provide functions such as drying, cleaning, blending, storing and merchandising grain. River elevators and the larger inland subterminals supply most of the corn destined for export markets. The figure below conveys the flow of United States corn destined for export markets.



B. IMPACT OF THE CORN MARKETING CHANNEL ON QUALITY

While the U.S. corn industry strives to prevent or minimize the loss of physical and sanitary quality as corn moves from the farm to export, there are points in the system where quality changes inevitably occur due to the biological nature of the grain. The following sections provide some insight into why corn quality may change as corn moves from the field to the vessel or railcar.

Drying and Conditioning

Farmers often harvest corn at moisture contents ranging from about 18 to 30%. This range of moisture contents exceeds safe storage levels, which are usually about 13 to 14%. Thus, wet corn at harvest must be dried to lower moisture content to become safe for storage and transport. Conditioning is the use of aeration fans to control temperatures and moisture content, both of which are important to monitor for storage stability. Drying and conditioning may occur either on a farm or at a commercial facility. When corn is dried, it can be dried by systems using natural air, low-temperature or high-temperature drying methods. High-temperature drying methods will often create more stress cracks in the corn and ultimately lead to more breakage during handling than natural air or low-temperature drying methods. However, high-temperature drying is often needed to facilitate the timely harvesting of grain.

Storage and Handling

In the United States, corn storage structures can be broadly categorized as upright metal bins, concrete silos, flat storage inside buildings or flat storage in on-ground piles. Upright bins and concrete silos with fully perforated floors or in-floor ducts are the most easily managed storage types, as they allow aeration with uniform airflow throughout the grain. Flat storage can be used for short-term storage. This occurs most often when corn production is higher than normal and surplus storage is needed. However, it is more difficult to install adequate aeration ducts in flat types of storage, and they often do not provide uniform aeration. In addition, on-ground piles are sometimes not covered and may be subjected to weather elements that can result in mold damage.

Handling equipment can involve vertical conveying by bucket elevators and/or horizontal conveying, usually by belt or en-masse conveyors. Regardless of how the corn is handled, some corn breakage will occur. The rate of breakage will vary by types of equipment used, the severity of the grain impacts, grain temperature, moisture content and by corn quality factors such as stress cracks or hardness of endosperm. As breakage levels increase, more fines (broken pieces of corn) are created, which leads to less uniformity in aeration and ultimately to a higher risk for fungal invasion and insect infestation.

Cleaning

Cleaning corn involves scalping or removing large non-corn material and sieving to remove small, shriveled kernels, broken pieces of kernels and fine material. This process reduces the amount of BCFM found in the corn. The potential for breakage and initial percentages of broken kernels, along with the desired grade factor, dictate the amount of cleaning needed to meet contract specifications. Cleaning can occur at any stage of the marketing channel where cleaning equipment is available.

Transporting Corn

The U.S. grain transportation system is arguably one of the most efficient in the world. It begins with farmers transporting their grain from the field to on-farm storage or commercial grain facilities using either large wagons or trucks. Corn is then transported by truck, rail or barge to its next destination. Once at export facilities, corn is loaded onto vessels or railcars.

Corn quality changes during shipment in much the same manner as it changes during storage. Causes of these changes include moisture variability (non-uniformity) and moisture migration due to temperature differences, high humidity and air temperature, fungal invasion and insect infestation. However, there are some factors affecting grain transportation that make quality control during transport more difficult than in fixed storage facilities. First, there are few modes of transport equipped with aeration and, as a result, corrective actions for heating and moisture migration cannot take place during transport. Another factor is the accumulation of fine material (spout-lines) beneath the loading spout when loading railcars, barges and vessels. This results in whole kernels tending to roll to the outer sides, while fine material segregates in the center. Similar segregation occurs during the unloading process at each step along the way to the final destination.

Implications on Quality

The intrinsic quality attributes, such as oil, protein and starch concentrations, remain essentially unchanged in a corn kernel between harvest and export, assuming negligible kernel respiration or mold damage. However, as corn moves through the U.S. corn marketing channel, corn from multiple sources is mixed together. As a result, the average for a given intrinsic quality characteristic is determined by the quality levels of the corn from multiple sources. Other changes occur in physical and sanitary quality characteristics. These include test weight, damaged kernels, broken kernels, stress crack levels, moisture content and variability, foreign material and mycotoxin levels.

C.U.S. GOVERNMENT INSPECTION AND GRADING

Purpose

Global corn supply chains need verifiable, predictable and consistent oversight measures that fit the diverse needs of all end-users. Oversight measures, implemented through standardized inspection procedures and grading standards, are established to provide:

- Information for buyers about the quality of grain at the time of loading for transport to the buyer, and
- Food and feed safety protection for the end-users.

The United States is recognized globally as having a combination of official grades and standards that are typically used for exporting grains and referenced in export contracts. U.S. corn sold by grade and shipped for foreign commerce must be officially inspected and weighed by FGIS or an official service provider delegated or designated by FGIS to do so (with a few exceptions). In addition, all corn exports must be tested for aflatoxins, unless the contract specifically waives this requirement. Qualified state and private inspection agencies are permitted to be designated by FGIS as official agents to inspect and weigh corn at specified interior locations. In addition, certain state inspection agencies can be delegated by FGIS to inspect and weigh grain officially at certain export facilities. Supervision of these agencies' operations and methodologies is performed by FGIS field office personnel.

Inspection and Sampling

The loading export elevator provides FGIS or the delegated state inspection agency a load order specifying the quality of the corn to be loaded as designated in the export contract. The load order specifies the U.S. grade, moisture content and all other requirements which have been agreed upon in the contract between the foreign buyer and the U.S. supplier, plus any special requirements requested by the buyer, such as minimum protein concentration, maximum moisture content or other special requirements. The official inspection personnel determine and certify that the corn loaded in the vessel or railcar meets the requirements of the load order. Independent laboratories can be used to test for quality factors not mandated to be performed by FGIS or for which FGIS does not have the local ability to test.

Shipments or “lots” of corn are divided into “sublots.” Representative samples for grading are obtained from these sublots using a diverter sampling device approved by FGIS. This device takes a primary portion approximately every 200 to 500 bushels (about 5.1 to 12.7 metric tons) from the moving grain stream just after the final elevation before loading into the vessel, shipping bin or railcar. The primary portions are usually further reduced by a secondary sampler, and incremental portions are combined by subplot and inspected by licensed inspectors. The results are entered into a log, and a statistical loading plan is applied to ensure not only that the average result for each factor meets the contract specifications, but also to ensure the lot is reasonably uniform in quality. Any subplot that does not meet uniformity criteria on any factor must be returned to the elevator or certified separately. The average of all subplot results for each factor is reported on the final official certificate. The FGIS sampling method provides a truly representative sample, while other commonly used methods may yield non-representative samples of a lot due to the uneven distribution of corn in a truck, railcar or in the hold of a vessel.

Grading

Yellow corn is divided into five U.S. numerical grades and U.S. Sample Grade. Each grade has limits for test weight, BCFM, total damaged kernels and heat-damaged kernels as a subset of total damage. The limits for each grade are summarized in the table shown in the “U.S. Corn Supporting Information” section of this report. In addition, FGIS provides certification of moisture content and aflatoxin results. Export contracts for corn can also specify other conditions or attributes related to the cargo, such as stress cracks, protein or oil concentrations and other mycotoxin results. In some cases, independent labs are used to conduct tests not required by FGIS.

Since the limits on all official grade factors (such as test weight and total damage) cannot always be met simultaneously, some grade factors may be better than the limit for a specified grade, but they cannot be worse. For that reason, most contracts are written as “U.S. No. 2 or better” or “U.S. No. 3 or better.” This permits some grade factor results to be at or near the limit for that grade, while other factor results are “better than” that grade.

A. OVERVIEW

The key points for the survey design and sampling and statistical analysis for this 2025/2026 *Export Cargo Report* are as follows:

- Following the methodology developed for the previous 14 *Export Cargo Reports*, samples were proportionately stratified according to ECAs – the Gulf, Pacific Northwest and Southern Rail.
- A total of 430 samples was targeted to achieve no more than a 10% Relative ME for the U.S. Aggregate level. These samples were to be collected from the ECAs as follows: 248 from the Gulf, 97 from the Pacific Northwest and 85 from the Southern Rail.
- A total of 430 samples was collected for this survey. The number of samples ultimately collected and tested from each ECA matched the number of samples targeted by ECA.
- Southern Rail ECA samples were provided by official agencies designated by FGIS that inspect and grade rail shipments of corn destined for export to Mexico. Gulf and Pacific Northwest samples were collected by FGIS field offices at ports in the respective ECAs.
- As in all previous *Export Cargo Reports*, chemical composition, physical factors and mycotoxin testing for the Gulf and Pacific Northwest ECAs was conducted on samples provided by FGIS based on our protocol for selection. Grade factor results from these two ECAs were obtained from FGIS's *Export Grain Report*, which contains grade factor data from corn shipments exported during the sampling period.
- To evaluate the statistical validity of the results, the Relative ME was calculated for each quality factor for the U.S. Aggregate and each of the three ECAs. The Relative ME for each of the quality factor results was not more than 10% at the U.S. Aggregate level. However, the Relative ME exceeded 10% in the Pacific Northwest and Southern Rail ECAs for total damage. It also exceeded 10% in the Southern Rail ECA for BCFM and stress cracks.
- Two-tailed t-tests at the 95% confidence level were calculated to measure statistical differences between this year's quality factor averages and those from this year's *Harvest Report*, the previous two *Export Cargo Reports*, the 5YA and the 10YA.

B. SURVEY DESIGN AND SAMPLING

Survey Design

For the *2025/2026 Export Cargo Report*, the target population was yellow commodity corn from the 12 key U.S. corn-producing states representing over 90% of the estimated 2025/2026 U.S. corn exports. A **proportionate stratified sampling** technique was used to ensure a sound statistical sampling of U.S. yellow corn exports. Two key characteristics define the sampling technique for this report: the **stratification** of the population to be sampled and the **sampling proportion** per subpopulation or stratum.

Stratification involves dividing the survey population of interest into subpopulations called strata. For the *Export Cargo Reports*, the key corn-exporting areas in the United States are divided into three geographical groupings, which we refer to as ECAs. These three ECAs are identified by the three major pathways to export markets:

- The Gulf ECA consists of areas that typically export corn through U.S. Gulf ports.
- The Pacific Northwest ECA includes areas that usually export corn through Pacific Northwest ports.
- The Southern Rail ECA comprises areas that generally export corn by rail to Mexico.


Using data from USDA, each ECA's proportion of the total expected annual yellow corn exports for the 2025/2026 corn marketing year was calculated. This average share of exports was used to determine the **sampling proportion** (the percent of total samples per ECA) and, ultimately, the number of yellow corn samples to be collected from each ECA.

The **number of samples** collected within each ECA was established so the Council could estimate the true U.S. Aggregate average for the various quality factors with a certain level of precision. The level of precision chosen for the *Export Cargo Report* was a Relative ME of not more than 10%, estimated at a 95% level of confidence.

To determine the number of samples for the targeted Relative ME, ideally, the population variance (i.e., the variability of the quality factor in the corn exports) for each of the quality factors should be used. The more variation among the levels or values of a quality factor, the more samples that are needed to estimate the true mean with a given confidence limit. In addition, the variances of the quality factors typically differ from one another. As a result, different sample sizes for each of the quality factors would be needed for the same level of precision.

Since the population variances for the 12 quality factors evaluated for this year's corn exports were not known, the variance estimates from previous editions of the *Export Cargo Report* were used as estimates of the population variance. The targeted number of samples for the desired level of precision for all quality factors were then calculated using these data.

Based on these historical data, 430 samples would allow the Council to estimate the true averages of the quality characteristics with the desired level of precision for the U.S. Aggregate. Applying the sampling proportions previously defined to the total of 430 samples resulted in the following number of targeted samples from each ECA as shown in the map below.



Beginning with the *2019/2020 Export Cargo Report*, a minimum of 180 samples was targeted to be tested for DON and horneous endosperm instead of the entire sample set. The same 180 samples tested for DON and horneous endosperm were also tested for fumonisin. The *2019/2020 Export Cargo Report* was the first *Export Cargo Report* in which this mycotoxin was tested. In terms of horneous endosperm, the Relative ME for this quality factor never exceeded 0.3% (well below the targeted level of 10.0%) in the samples tested from the first eight previous reports. Thus, reducing the number of samples tested for horneous endosperm would likely keep the precision of this quality factor's estimates well below the targeted level of 10.0%. In the *2019/2020 Export Cargo Report*, the first *Export Cargo Report* in which only 180 samples were tested for horneous endosperm, the Relative ME for this quality factor was only 0.4%.

Beginning with the *2020/2021 Export Cargo Report*, the target of 180 samples was extended to 100-kernel weight, kernel volume and kernel true density. Given the historical data collected from the first nine *Export Cargo Reports*, reducing the number of samples tested for these three additional quality factors would likely keep the precision of their estimates well below the targeted level of 10.0%.

Beginning with the *2021/2022 Export Cargo Report*, the same samples to be tested for aflatoxin, DON and fumonisin were also tested for ochratoxin A, trichothecenes (T-2) and zearalenone.


In the first eight years of the *Export Cargo Report*, the stress crack index was reported in addition to the percent stress cracks to indicate the severity of stress cracking. The stress crack index is determined using the following calculation:

$$[\text{SSC} \times 1] + [\text{DSC} \times 3] + [\text{MSC} \times 5]$$

Where

- SSC is the percentage of kernels with only one crack;
- DSC is the percentage of kernels with exactly two cracks; and
- MSC is the percentage of kernels with more than two cracks.

The U.S. Aggregate percent stress cracks and stress crack index from the first eight *Export Cargo Reports* is displayed in the scatter chart below. Given its strong correlation ($r = 0.97$) to percent stress cracks, it was determined that the stress crack index provided limited additional value and was discontinued following the *2018/2019 Export Cargo Report*.



Sampling

The sampling was administered by FGIS and participating official service providers as part of their inspection services. FGIS sent instruction letters to the Gulf and Pacific Northwest field offices and to the domestic inspection offices. The sampling period began in November 2025 and concluded in early-February 2026. The FGIS field offices in the respective ECAs responsible for overseeing the sample collection within their region were as follows: Gulf – New Orleans, Louisiana; Pacific Northwest – Olympia, Washington (Washington State Department of Agriculture); and Southern Rail – FGIS Domestic Inspection Operations Office in Kansas City, Missouri.

While the sampling process is continuous throughout the loading of an ocean-bound vessel, a shipment or “lot” of corn is divided into “sublots” for the purpose of determining the uniformity of quality. Sublot size is based on the hourly loading rate of the elevator and the capacity of the vessel being loaded. Sublot sizes range from 30,000 to 120,000 bushels. All sublot samples are inspected.

Representative sublot samples from the ports in the Gulf and Pacific Northwest ECAs were collected as ships were loaded. Samples for grading are obtained by a diverter sampler approved by FGIS. The diverter sampler “cuts” (or diverts) a representative portion from a flowing stream of grain at periodic intervals. A cut occurs every few seconds, or about every 200 to 500 bushels (about 5.1 to 12.7 metric tons), as the grain is being assembled for export. The frequency is regulated by an electronic timer controlled by official inspection personnel who periodically determine that the mechanical sampler is functioning properly. For each sample, a minimum of 2,700 grams was provided by the FGIS field staff and the Washington State Department of Agriculture.

As in all previous *Export Cargo Reports*, samples were provided by FGIS from the Gulf and Pacific Northwest ECAs and used for chemical composition, physical factors and mycotoxin testing. However, grade factor results from these two ECAs were obtained from FGIS’s Export Grain Report instead of from our testing the samples provided by FGIS. This *2025/2026 Export Cargo Report* is the fourth year in which the results presented reflect this change in protocol. To allow for comparisons across years, grade factor data from all previous reports were replaced with data obtained from the FGIS Export Grain Report. Annual inspection data are available to the public at <https://fgisonline.ams.usda.gov/ExportGrainReport/default.aspx>.

The FGIS Export Grain Report provides shipment-level grade factor and moisture data, the contract grade specified by the trading partners, the amount shipped in metric tons and the inspection date. These data were filtered by date to correspond with the same timeframe from which the samples provided by FGIS were collected. Complete grade factor data are available from nearly all vessels loaded from the Gulf and Pacific Northwest ECAs, which provides the best possible representation of the average grade factors loaded from these ECAs during the sampling period since all vessels loaded with corn are sampled with only a few exceptions. These exceptions include the following:

- The dataset was filtered to exclude rail, truck, and containerized exports. Only vessel shipments were retained to best match the population from which the samples provided by FGIS were collected.
- Observations were scrubbed to exclude unrealistic values. The most common unrealistic value was a number entered into the incorrect field such as a test weight value entered into the total damage field. Only a small fraction of observations contained errors, and the few observations containing unrealistic values were excluded from the dataset.
- Several types of exports are exempt from FGIS inspection. The reasons for exemption include corn sold for seed, corn not sold by grade, corn exported by small exporters, specialty shipments of corn and grain exported by rail or truck to Canada and Mexico.

While the dataset contains shipment information from the Southern Rail ECA, this *Export Cargo Report* continues to rely on the grade factor results collected from the samples provided by FGIS.¹ Grain exported by rail to Mexico is exempt from FGIS inspection. While this dataset generally includes corn exported to Mexico by rail or truck, grade factor data are often incomplete for these shipments.

For the Southern Rail ECA, representative samples were taken about every 200 bushels (5.1 metric tons) at domestic interior elevators using a diverter sampler to ensure uniform sampling. Only trains of yellow corn inspected for export to Mexico were sampled. The Southern Rail ECA official service providers submitted file samples exclusively. These samples were collected and tested for grade factors and aflatoxin at the time of sampling and then kept on file at the official service providers to be retested in the case of disputes. Each file sample weighed approximately 1,000 grams and represented a composite of five railcars. File samples were mailed to the Illinois Crop Improvement Association's Identity Preserved Grain Laboratory (IPG Lab) after their retention dates were reached, which were generally 30 days after loading. Grade factor tests for the Southern Rail samples were conducted by Champaign-Danville Grain Inspection (CDGI) in Urbana, Illinois. CDGI is the official grain inspection service provider for East Central Illinois as designated by USDA FGIS. Of the 85 samples received from the Southern Rail ECA, 28 weighed less than 1,000 grams. Grade factor results were not possible as 1,000 grams are necessary to conduct these tests.

¹Moisture content is the one exception to this protocol. While often incomplete, FGIS data from the Southern Rail ECA was deemed to offer a better representation of moisture content compared to the moisture content determined from the physical samples collected as there was significant potential for samples to decrease in moisture during storage and transit.

C. STATISTICAL ANALYSIS

The sample test results for grade factors, chemical composition and physical factors were summarized for the U.S. Aggregate and also for the three ECAs (Gulf, Pacific Northwest and Southern Rail). Summarized grade factor data are also reported by the following two contracted grade categories²:

- Shipments contracted to be loaded as “U.S. No. 2” or better
- Shipments contracted to be loaded as “U.S. No. 3” or better

Within this *2025/2026 Export Cargo Report* is a simple average of the quality factors’ averages and standard deviations of the previous five *Export Cargo Reports* (2020/2021, 2021/2022, 2022/2023, 2023/2024 and 2024/2025). These simple averages are calculated for the U.S. Aggregate and each of the three ECAs and are referred to as the “5YA” in the text and summary tables of the report. References to the “10YA” are also made throughout the report. The 10YA represents the simple average of the quality factors’ averages from the *2015/2016 Export Cargo Report* through the *2024/2025 Export Cargo Report*.

The Relative ME was calculated for each of the quality factors tested for this study at the U.S. Aggregate level and for each of the ECAs. The Relative ME was not more than 10% for all the quality attributes at the U.S. Aggregate level. However, it exceeded 10% in the Pacific Northwest ECA for total damage (12%) and in the Southern Rail ECA for stress cracks (20%), total damage (22%) and BCFM (14%). While the level of precision for these estimates is less than desired, the levels of Relative ME do not invalidate the estimates. The averages for the quality factors are the best possible unbiased estimates of the true population means. However, they are estimated with greater uncertainty than the quality factors with a Relative ME of less than 10%. Footnotes in the summary tables for “Grade Factors” and “Physical Factors” indicate the attributes for which the Relative ME exceeded 10%.

References in the “Quality Test Results” section to statistical differences were validated by 2-tailed t-tests at the 95% confidence level. These tests were calculated to determine statistical differences between quality factor averages from this *Export Cargo Report* and the following:

- This year’s *Harvest Report* and
- Each of the previous two *Export Cargo Reports*, the 5YA and the 10YA.

²Grade classifications for Gulf and Pacific Northwest samples were determined by the contracted grade agreed to by the transacting parties rather than the actual grade of the corn at loading. Grade classifications for Southern Rail samples were determined from the actual grade of the corn because contracted grade of the shipments sampled is unknown. U.S. Aggregate results for grades other than U.S. No. 2 or U.S. No. 3 are not reported due to limited number of samples. Chemical composition and physical factor results were not summarized by grade classification because the contracted grade of the physical samples collected from the Gulf and Pacific Northwest shipments is unknown.

The 2025/2026 *Export Cargo Report* samples were sent directly from the FGIS field offices and official service providers to the IPG Lab in Champaign, Illinois. IPG Lab conducted chemical composition, physical factor and mycotoxin testing. Grade factor tests on samples collected from the Southern Rail ECA were conducted by Champaign-Danville Grain Inspection (CDGI) in Urbana, Illinois.¹ CDGI is the official grain inspection service provider for East Central Illinois as designated by USDA FGIS. The grade testing procedures were in accordance with FGIS's *Grain Inspection Handbook* and are described in the following section. IPG Lab has received accreditation under the ISO/IEC 17025:2017 International Standard for many of the tests. The full scope of accreditation is available at <https://www.ilcrop.com/labservices>.

A. GRADE FACTORS

Test Weight

Test weight is a measure of the volume of grain required to fill a Winchester bushel (2,150.42 cubic inches). Test weight is a part of the FGIS Official U.S. Standards for Corn grading criteria.

The test involves filling a test cup of known volume through a funnel held at a specific height above the test cup to the point where grain begins to pour over the test cup's sides. A strike-off stick is used to level the grain in the test cup, and the grain remaining in the cup is weighed. The weight is then converted to and reported in the traditional U.S. unit, pounds per bushel (lb/bu).

Broken Corn and Foreign Material

BCFM is part of the FGIS Official U.S. Standards for Grain and grading criteria.

The BCFM test determines the amount of all matter that passes through a 12/64th-inch round-hole sieve and all matter other than corn that remains on the top of the sieve. BCFM measurement can be separated into broken corn and foreign material. Broken corn is defined as all material passing through a 12/64th-inch round-hole sieve and retained on a 6/64th-inch round-hole sieve. The definition of foreign material is all material passing through the 6/64th-inch round-hole sieve and the coarse non-corn material retained on top of the 12/64th-inch round-hole sieve. While FGIS can report broken corn and foreign material separately if requested, BCFM is the default measurement and is provided for the *Export Cargo Report*. BCFM is reported as a percentage of the initial sample by weight.

¹Grade factor tests on samples collected from the Gulf and Pacific Northwest ECAs were conducted by FGIS during the initial inspection of the vessel. Average shipment-level results are made available by FGIS at <https://fgisonline.ams.usda.gov/ExportGrainReport/default.aspx>.

Total Damage and Heat Damage

Total damage is part of the FGIS Official U.S. Standards for Grain grading criteria.

A trained and licensed inspector visually examines a representative working sample of 250 grams of BCFM-free corn for damaged kernels. Types of damage include blue-eye mold, cob rot, dryer-damaged kernels (different from heat-damaged kernels), germ-damaged kernels, heat-damaged kernels, insect-bored kernels, mold-damaged kernels, mold-like substance, silk-cut kernels, surface mold (blight), mold (pink *Epicoccum*) and sprout-damaged kernels. Total damage is reported as the weight percentage of the working sample that is total damaged grain.

Heat damage is a subset of total damage and consists of kernels and pieces of corn kernels that are materially discolored and damaged by heat. Heat-damaged kernels are determined by a trained and licensed inspector visually inspecting a 250-gram sample of BCFM-free corn. Heat damage, if found, is reported separately from total damage.

B. CHEMICAL COMPOSITION

Near-Infrared Transmission Spectroscopy (NIR) Proximate Analysis

The chemical composition (protein, oil and starch concentrations) of corn is measured using NIR. The technology uses unique interactions of specific wavelengths of light with each sample. It is calibrated to traditional chemistry methods to predict protein, oil and starch concentrations in the sample. This procedure is nondestructive to the corn.

Chemical composition tests for protein, oil and starch were conducted using a 550 to 600-gram sample in a whole-kernel Foss Infratec 1241 NIR instrument. The NIR was calibrated to chemical tests, and the standard errors of predictions for protein, oil and starch were about 0.22%, 0.26% and 0.65%, respectively. Comparisons of the Foss Infratec 1229 used in *Export Cargo Reports* before 2016 to the Foss Infratec 1241 on 21 laboratory check samples showed the instruments averaged within 0.25%, 0.26% and 0.25% points of each other for protein, oil and starch, respectively. Results are reported on a dry basis percentage (percent of non-water material).

C. PHYSICAL FACTORS

100-Kernel Weight, Kernel Volume and Kernel True Density

The 100-kernel weight is determined from the average weight of two 100-kernel replicates using an analytical balance that measures to the nearest 0.1 milligrams. The averaged 100-kernel weight is reported in grams.

The kernel volume for each 100-kernel replicate is calculated using a helium pycnometer and is expressed in cubic centimeters (cm³) per kernel. Kernel volumes usually range from 0.14 cubic centimeters to 0.36 cubic centimeters per kernel for small and large kernels, respectively.

True density of each 100-kernel sample is calculated by dividing the mass (or weight) of the 100 externally sound kernels by the volume (displacement) of the same 100 kernels. The two replicate results are averaged. True density is reported in grams per cubic centimeter (g/cm³). True densities typically range from 1.20 grams per cubic centimeter to 1.30 grams per cubic centimeter at “as is” moisture contents of about 12 to 15%.

Stress Crack Analysis

Stress cracks are evaluated by using a backlit viewing board to accentuate the cracks. A sample of 100 intact kernels with no external damage is examined kernel by kernel. The light passes through the horneous or hard endosperm, so each kernel’s stress crack damage can be evaluated. Kernels are sorted into two categories: (1) no cracks; (2) one or more cracks. Stress cracks, expressed as a percent, are all kernels containing one or more cracks divided by 100 kernels. Lower levels of stress cracks are always better since higher stress cracks lead to more breakage in handling. Some end-users will specify by contract the acceptable level of cracks based on the intended use.

Whole Kernels

In the whole kernels test, 50 grams of cleaned (BCFM-free) corn are inspected kernel by kernel. Kernels that are cracked, broken, chipped or showed significant pericarp damage are removed. The whole kernels are then weighed, and the result is reported as a percentage of the original 50-gram sample. Some companies perform the same test but report the “cracked & broken” percentage. A whole kernel score of 97.0% equates to a cracked & broken rating of 3.0%.

Horneous (Hard) Endosperm

The horneous (or hard) endosperm test is performed by visually rating 20 externally sound kernels placed germ facing up on a backlit viewing board. Each kernel is rated for the estimated portion of the kernel's total endosperm that is horneous endosperm. The soft endosperm is opaque and will block light, while the horneous endosperm is translucent. The rating is made from standard guidelines based on the degree to which the soft endosperm at the crown of the kernel extends down toward the germ. The average of horneous endosperm ratings for the 20 externally sound kernels is reported. Ratings of horneous endosperm are made on a scale of 70 to 100%, though most individual kernels fall in the 70 to 90% range.

D. MYCOTOXINS

To report the frequency of occurrences of aflatoxin, DON and fumonisin for the *2025/2026 Export Cargo Report*, IPG Lab performed the mycotoxin testing using test kits approved by FGIS. For this study, a 1,000-gram laboratory sample was subdivided from the survey sample of shelled kernels for the mycotoxin analysis. The one-kilogram survey sample was ground in a Retsch ZM 200 mill, with a distance ring sieve size of 1 millimeter and trapezoidal holes so that 60 to 75% would pass through a 20-mesh screen. From this well-mixed ground material, a 50-gram test portion was removed for each mycotoxin tested. EnviroLogix AQ 309 BG, AQ 304 BG and AQ 411 BG quantitative test kits were used for the aflatoxin, DON and fumonisin analysis, respectively. DON and fumonisin were extracted with water (5:1), while the aflatoxin was extracted with buffered water (3:1). The extracts were tested using the EnviroLogix QuickTox lateral flow strips, and the QuickScan system quantified the mycotoxins.

The EnviroLogix quantitative test kits report specific concentration levels of the mycotoxin if the concentration level exceeds a specific level called a "Limit of Detection." The limit of detection is defined as the lowest concentration level that can be measured with an analytical method that is statistically different from measuring an analytical blank (absence of a mycotoxin). The limit of detection will vary among different types of mycotoxins, test kits and commodity combinations. The limit of detection for the EnviroLogix AQ 309 BG is 2.7 parts per billion for aflatoxin. The limit of detection for DON using the EnviroLogix AQ 304 BG is 0.1 parts per million. For the fumonisin tests, the EnviroLogix AQ 411 BG has a limit of detection of 0.1 parts per million. FGIS has issued a letter of performance for the quantification of aflatoxin, DON and fumonisin using the Envirologix AQ 309 BG, AQ 304 BG and AQ 411 BG kits, respectively.



In the first nine *Export Cargo Reports*, FGIS provided aflatoxin results. Official FGIS protocol for aflatoxin testing requires the grinding of at least 10 pounds of shelled corn from vessel subplot and composite inspections. Beginning with the *2020/2021 Export Cargo Report*, IPG Lab received a minimum of 1,000 grams for aflatoxin testing. This represented a change in protocol from the first nine *Export Cargo Reports*.

According to the official FGIS aflatoxin testing protocol utilized in the first nine *Export Cargo Reports*, a 10-pound sample is ground using a FGIS-approved grinder. Following the grinding stage, two 500-gram ground portions are removed from the 10-pound comminuted sample using a riffle divider. From one of the 500-gram ground portions, a 50-gram test portion is randomly selected for testing. After adding the proper extraction solvent to the 50-gram test portion, aflatoxins are quantified. The following FGIS-approved quantitative test kits may have been used: Charm Sciences, Inc. ROSA® FAST, WET-S3 or WET-S5 Aflatoxin Quantitative Tests; EnviroLogix, Inc. QuickTox™ Kit for QuickScan Aflatoxin Flex AQ 309 BG; Hygienea LLC Mycotox Total Aflatoxin ELISA; Neogen Corporation Reveal Q+ MAX for Aflatoxin, Reveal Q+ for Aflatoxin, or Veratox® Aflatoxin Quantitative Test (8030 or 8035); R-Biopharm, Inc. RIDASCREEN® FAST Aflatoxin ECO; Romer Labs, Inc. FluoroQuant Afla or AgraStrip Total Aflatoxin Quantitative Test WATEX; PerkinElmer Inc. AuroFlow AQ Afla Strip Test; or VICAM AflaTest™ or Afla-V AQUA.

The *Export Cargo Report* also began testing samples for ochratoxin A, T-2 and zearalenone beginning with the *2021/2022 Export Cargo Report*. The testing for these three additional mycotoxins complements the information provided by the test results from the two mycotoxins tested since the *2011/2012 Export Cargo Report* (aflatoxin and DON) and fumonisin (tested since the *2019/2020 Export Cargo Report*). EnviroLogix AQ 113 BG, AQ 314 BG, and AQ 412 BG quantitative test kits were used for ochratoxin A, T-2 and zearalenone, respectively. The EnviroLogix AQ 113 BG quantitative test kit used for the ochratoxin A tests has a limit of detection of 1.5 parts per billion. The ochratoxin A was extracted with a grain buffer (five milliliters per gram). For the T-2 tests, the AQ 314 BG quantitative test kit has a limit of detection of 50 parts per billion. T-2 was extracted with water (five milliliters per gram). The EnviroLogix AQ 412 BG quantitative test kit used for the zearalenone tests has a limit of detection of 50 parts per billion. The zearalenone test uses a 25-gram test portion of corn. The zearalenone was extracted using a reagent of EB17 extraction powder and a water buffer of 75 milliliters per sample.



GRADE FACTORS AGGREGATE HARVEST AND EXPORT CARGO COMPARISON

Since 2011, the U.S. Grains & BioProducts Council's *Export Cargo Reports* have provided clear, concise and consistent information about the quality of each U.S. crop entering international merchandising channels. This series of quality reports has used consistent and transparent methodology to allow for insightful comparisons across time. The following charts display the average U.S. Aggregate from all *Harvest Reports* and *Export Cargo Reports* for each quality factor tested to provide historical context to this year's results.








**GRADE FACTORS
AGGREGATE HARVEST AND EXPORT CARGO COMPARISON**






CHEMICAL COMPOSITION AGGREGATE HARVEST AND EXPORT CARGO COMPARISON








**PHYSICAL FACTORS
AGGREGATE HARVEST AND EXPORT CARGO COMPARISON**





PHYSICAL FACTORS AGGREGATE HARVEST AND EXPORT CARGO COMPARISON









MYCOTOXINS
HARVEST AND EXPORT CARGO COMPARISON



MYCOTOXINS HARVEST AND EXPORT CARGO COMPARISON





U.S. CORN GRADES AND GRADE REQUIREMENTS

Grade	Minimum Test Weight per Bushel (Pounds)	Maximum Limits of		
		Damaged Kernels		Broken Corn and Foreign Material (Percent)
		Heat Damaged (Percent)	Total (Percent)	
U.S. No. 1	56.0	0.1	3.0	2.0
U.S. No. 2	54.0	0.2	5.0	3.0
U.S. No. 3	52.0	0.5	7.0	4.0
U.S. No. 4	49.0	1.0	10.0	5.0
U.S. No. 5	46.0	3.0	15.0	7.0

U.S. Sample Grade is corn that: (a) Does not meet the requirements for the grades U.S. Nos. 1, 2, 3, 4, or 5; or (b) Contains stones with an aggregate weight in excess of 0.1% of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria spp.*), 2 or more castor beans (*Ricinus communis L.*), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cockleburrs (*Xanthium spp.*), or similar seeds singly or in combination, or animal filth in excess of 0.2% in 1,000 grams; or (c) Has a musty, sour, or commercially objectionable foreign odor; or (d) Is heating or otherwise of distinctly low quality.

Source: Code of Federal Regulations, Title 7, Part 810, Subpart D, United States Standards for Corn

U.S. AND METRIC CONVERSIONS

Corn Equivalents	Metric Equivalents
1 bushel = 56 pounds (25.40 kilograms)	1 pound = 0.4536 kilograms
39.368 bushels = 1 metric ton	1 hundredweight = 100 pounds or 45.36 kilograms
15.93 bushels/acre = 1 metric ton/hectare	1 metric ton = 2204.6 pounds
1 bushel/acre = 62.77 kilograms/hectare	1 metric ton = 1000 kilograms
1 bushel/acre = 0.6277 quintals/hectare	1 metric ton = 10 quintals
56 pounds/bushel = 72.08 kilograms/hectoliter	1 quintal = 100 kilograms
	1 hectare = 2.47 acres

ABBREVIATIONS

cm ³ = cubic centimeters
g = grams
g/cm ³ = grams per cubic centimeter
kg/hl = kilograms per hectoliter
lb/bu = pounds per bushel
ppb = parts per billion
ppm = parts per million



A global network of professionals building worldwide demand and developing markets for U.S. grains and ethanol.



HEADQUARTERS

20 F Street NW, Suite 900 • Washington, D.C., 20001
 Phone: 202-789-0789 • Fax: 202-898-0522
 Email: grains@grains.org • Website: grains.org

PEOPLE'S REPUBLIC OF CHINA: Beijing

Tel1: 011-86-10-6505-1314 • Tel2: 011-86-10-6505-2320
 Fax: 011-86-10-6505-0236 • china@grains.org

JAPAN: Tokyo

Tel: 011-81-3-6206-1041 • Fax: 011-81-3-6205-4960
japan@grains.org • www.grainsjp.org

KOREA: Seoul

Tel: 011-82-2-720-1891 • Fax: 011-82-2-720-9008
seoul@grains.org

TAIWAN: Taipei

Tel: 011-886-2-2523-8801 • Fax: 011-886-2-2523-0149
taipei@grains.org

MIDDLE EAST, AFRICA AND EUROPE: Tunis

Tel: 011-216-71-191-640 • Fax: 011-216-71-191-650
tunis@grains.org

INDIA: New Delhi

Tel: +91-11-4603-6437 • southasia@grains.org

SOUTH EAST ASIA: Kuala Lumpur

Tel: 011-603-2789-3288 • sea-oceania@grains.org

LATIN AMERICA: Panama City

Tel: 011-507-315-1008 • lta@grains.org

MEXICO: Mexico City

Tel1: 011-52-55-5282-0244 • Tel2: 011-52-55-5282-0973
 Tel3: 011-52-55-5282-0977 • mexico@grains.org